

APPENDIX T

Specific Management Recommendations for Endangered Species

The following information and management guidelines are from the 130 page book "Endangered and Threatened Animals of Texas - Their Life History and Management", by Linda Campbell. Published by the Texas Parks and Wildlife Press, Austin, Texas in 1995. Distributed by the University of Texas Press, Austin, Texas, and revised in 2003 as an electronic book available on the TPWD website at www.tpwd.state.tx.us.

Black-capped Vireo



Scientific Name: *Vireo atricapillus*
Federal Status: Endangered, 10/6/87 • State
Status: Endangered

Description

The Black-capped Vireo is a 4.5 inch insect-eating songbird. Mature males are olive green above and white below with faint greenish-yellow flanks. The crown and back of the head is black with a partial white eyering. The iris is brownish-red and the bill black. The plumage on the back of the female is duller than the male. Females have a medium to

dark gray head with a blackish ring around the white surrounding the eye (this generally distinguishes the female from the second year male).

Distribution and Habitat

Historical records from 1852-1956 show that the Black-capped Vireo once occurred and nested from central Kansas, Oklahoma, Texas and into northern Mexico. Today, Blackcapped Vireos are known to nest in central and southwest Texas, a few counties in central Oklahoma, and in Coahuila and Nuevo Leon, Mexico, although less is known of their status in Mexico. Black-capped Vireos winter along the western coast of Mexico.

The descriptions of habitat presented in this document are intended to help landowners determine if they have Black-capped Vireo habitat on their property. Not all sites within the habitat types described will be used by Black-capped Vireos. It is only where individuals of this species occupy the identified habitat types during the breeding season that special management considerations such as those provided in these guidelines need to be considered. In Texas, vireo habitat is found on rocky limestone soils of the Edwards Plateau, Cross Timbers and Prairies, eastern Trans-Pecos and, to a limited extent, on igneous soils in the Chisos Mountains. Although Blackcapped Vireo habitat throughout Texas is highly variable with regard to plant species, soils, temperature, and rainfall, all habitat types are similar in vegetation structure; i.e. the "overall look" is somewhat similar although the plant species vary.

Vireos require broadleaf shrub vegetation reaching to ground level for nesting cover. They typically nest in shrublands and open woodlands with a distinctive patchy structure. Typical habitat is characterized by shrub vegetation extending from the ground to about 6 feet or more and covering about 30-60% or greater of the total area. In the eastern portion of the vireo's range, the shrub layer is often combined with an open, sparse to moderate tree canopy.

Patches of open grass or bare rock separate the clumps of shrubs and trees. In central Texas, this habitat is often regrowth from disturbances such as clearing, fire, and browsing. In the Edwards Plateau and Cross Timbers Regions, vireo habitat occurs where soils, topography, and land use produce scattered hardwoods with abundant low cover. Common broadleaved plants in vireo habitat in these regions include: Texas (Spanish) oak, Lacey oak, shin oak, Durand (scaleybark) oak, live oak, mountain laurel, evergreen sumac, skunkbush sumac, flameleaf sumac, redbud, Texas persimmon, Mexican buckeye, elbowbush and agarita. Although Ashe juniper is often part of the plant composition in vireo habitat, preferred areas usually have a low density and cover of juniper.

In the western Edwards Plateau and Trans-Pecos Regions, on the western edge of the vireo's range, the birds are often found in canyon bottoms and slopes where sufficient moisture is available to support diverse shrub vegetation. Dominant woody plants in this habitat type include sandpaper oak, Vasey oak, Texas kidneywood, Mexican walnut, Texas persimmon, lotebush, brasil, wafer ash, mountain laurel, cenizo, whitebrush, and guajillo. For all habitat types, the plant composition appears to be less important than the presence of adequate broad-leaved shrubs, foliage to ground level, and mixture of open grassland and woody cover. Deciduous and broad-leaved shrubs and trees throughout the vireo's range are also important in providing habitat for insects on which the vireo feeds.

Life History

Black-capped Vireos arrive in Texas from mid-March to mid-April. Adult males often arrive before females and first-year males to select their territories. Vireos' territories are often clustered in patches of suitable habitat. Although territories range in size from 1 to 16 acres, most territories are 5 to 10 acres. Males sing to attract mates and defend territories. Many males can be heard singing throughout the breeding season, but singing begins to decline by July. The vireo's song is described as hurried and harsh, composed of numerous phrases separated from one another by pauses of 1 to 3 seconds. Nesting begins after the females arrive in late March to early April. Both the male and female select the nest site and build the nest, but the female often completes it. First nests are built in about 6 to 9 days, but subsequent nests can be built in one day. The cup-shaped nest is suspended from its rim in a fork of a branch about 1 to 6 feet above the ground. However, most Black-capped Vireos nest at about "door-knob" height. Nests have been found in a variety of species including shin oak, scaleybark oak, Texas oak, Vasey oak, sumac, Texas persimmon, juniper, Texas redbud, Mexican buckeye and Texas mountain laurel. The vireo usually nests more than once in the same year. A new nest is constructed each time. Three to four eggs are usually laid in the first nesting attempt, but later clutches may contain only 2 to 3 eggs. The first egg is usually laid one day after completion of the nest, with one egg being laid each subsequent day. Incubation takes 14 to 17 days, and is shared by the male and female. Vireo chicks are fed insects by both adults. The young leave the nest 10 to 12 days after hatching. Fledglings are cared for by the female alone, the male alone, or by both adults. Sometimes the parents split the brood and each care for one or more young. Occasionally, males or females will leave the care of the young to their mate, and attempt another nesting effort. Vireos may live for more than five years, and usually return year after year to the same territory, or one nearby. The birds migrate to their wintering grounds on Mexico's western coast beginning in July, and are gone from Texas by mid-September.

Threats and Reasons for Decline

The Black-capped Vireo is vulnerable to changes in the abundance and quality of its habitat. Habitat may become unsuitable for vireos because of natural plant succession, sustained brood parasitism by the Brown-headed Cowbird, or because of human activities. Factors that can

adversely affect vireo habitat include broad-scale or improper brush clearing, fire suppression, over browsing by deer and livestock, and urbanization. Loss of tropical wintering habitat is also a concern, but requires further study. Poorly planned brush management practices on rangeland may remove too much low growing woody cover, especially when large acreages are treated at one time. This eliminates or reduces habitat value for vireos and for other wildlife, such as White-tailed deer, quail, small mammals, and various songbirds. Over browsing of broad-leaved shrubs by goats, deer, and exotic animals reduces the vegetation in the 2- to 4-foot zone, making it unsuitable for vireo nesting. Continued overuse of these preferred browse plants over many years may eventually eliminate them from the plant community, thus permanently altering the habitat.

In the absence of natural processes, active, well-planned land management is often required to maintain good vireo habitat, especially in the eastern portion of its range. Disturbance, particularly fire, plays an important role in maintaining, improving, or creating vireo habitat. The rangelands of central Texas, and the various plant communities these lands support, evolved under the influence of periodic fires. Historically, these natural and manmade fires maintained a matrix of open grassland, shrubland and woodland. Fire stimulated shrubs to sprout multiple stems at the base, thus providing areas of dense foliage at the 2- to 4-foot level, required by vireos. In the past, fire was responsible for maintaining or periodically returning some areas to vireo habitat. Today, prescribed burning, a valuable range and wildlife management tool occurs on many ranches throughout Texas. However, the combination of overgrazing, brush clearing, and lack of fire in the recent past has reduced vireo habitat in many other areas. Natural plant succession is less of a concern in the western portion of its range where suitable habitat persists for long periods.

Human activities have provided favorable habitat for the Brownheaded Cowbird, which parasitizes vireo nests. The cowbird is usually associated with livestock, farms, dairies, and grain fields, where it benefits from waste grain and insects. They may also be attracted to backyard bird feeders, trash dumps, or other urban areas where food and water are available. Cowbirds lay their eggs in other birds' nests, leaving the host bird to raise their young. The female cowbird often removes an egg or a nestling from the host nest before she lays an egg in it. Cowbird chicks hatch earlier than most hosts' young and are thus able to out-compete the smaller vireo nestlings for food and, consequently, the young vireos typically starve. While some birds remove cowbird eggs from their nest, the vireo does not, although it is known to abandon parasitized nests. Thus parasitized nests usually fail to produce vireos. The amount of brood parasitism varies greatly from one population to another throughout the state, ranging from 10 to over 90% of the nests. Brown-headed Cowbirds are also known to remove vireo chicks from active nests. Evidence indicates that sustained parasitism pressure may lead to local extinctions of vireo populations. Direct habitat loss and fragmentation due to urban and suburban development is a major threat in expanding urban areas of Travis, McLennan, Dallas, Bexar, and Kerr counties. Problems associated with suburban expansion, such as increases in predation by dogs, cats, raccoons, skunks, and jays, have also impacted the vireo.

Recovery Efforts

Research is underway to better understand the distribution, life history, habitat requirements, and land management practices affecting the Blackcapped Vireo. Population surveys during the breeding season are being conducted in known and potential habitat areas. Efforts to provide information and educational opportunities to landowners and the public regarding life history and habitat requirements of the vireo are also a vital part of the recovery effort. Major research and/or recovery efforts are being conducted on Department of Defense's Fort Hood and Camp Bullis, Travis County and the City of Austin's Balcones Canyonlands Preserve, the U.S. Fish

and Wildlife Services' Balcones Canyonlands National Wildlife Refuge, TPWD's Kerr Wildlife Management Area, properties owned and/or managed by The Nature Conservancy of Texas, and in Mexico. Additionally, Environmental Defense through their Safe Harbor Agreement with the U.S. Fish and Wildlife Service is assisting many landowners with thousands of acres to manage and/or create habitat for the vireo. Research is ongoing regarding the impact of cowbirds on vireo populations in Texas. Research efforts in Mexico are also underway to gather information concerning life history, habitat requirements, and conservation threats on the wintering range. TPWD biologists are monitoring populations on both state and private lands, and voluntary cowbird trapping is being conducted by more than 400 landowners in counties throughout the range of the vireo. Habitat conservation planning is underway in counties such as Travis and Bexar to allow for urban expansion and development while still conserving endangered species habitat. Intensive monitoring of a large population at the U.S. Army Fort Hood Military Installation is on-going. Finally, efforts to provide information, technical assistance, and incentives for private landowners to incorporate management for Black-capped Vireos into their livestock and wildlife operations are an essential part of the recovery process.

Where To See the Black-capped Vireo

A number of state lands offer opportunities to see and learn more about the Black-capped Vireo. These include Colorado Bend State Park State Park (SP), Devils River State Natural Area (SNA), Kerr Wildlife Management Area, Kickapoo Cavern SP, Lost Maples SNA, and Hill Country SNA. Also, the Balcones Canyonlands National Wildlife Refuge near Austin offers additional opportunities to see Black-capped Vireos. Because the Black-capped Vireo is an endangered species, birders and other observers should carefully follow certain viewing ethics. Observers should be careful not to flush birds from the nest or disturb nests or young. Black-capped Vireos should be viewed only from a distance with binoculars. Do not use recorded calls of the Black-capped Vireo or the Screech Owl to attract birds, and be careful that your presence does not unduly disturb or stress the birds.

How You Can Help

You can help by learning more about the habitat requirements of the Blackcapped Vireo and incorporating management practices which create or maintain habitat for these birds. You can also encourage and support private landowners who are managing their land to protect and provide habitat for endangered species. The Black-capped Vireo is a beautiful songbird and is much sought after among people who enjoy birdwatching and nature study. Possibilities exist for landowners to take advantage of the growing demand for natural history tours and vacations. Landowners interested in more information concerning nature-based tourism opportunities should contact the Wildlife Diversity Branch, Texas Parks and Wildlife Department, Austin (800) 792-1112; Environmental Defense, Austin (512) 478-5161; the Nature Conservancy, San Antonio (210) 224-8774. You can also be involved with the conservation of Texas' nongame wildlife resources by supporting the Special Nongame and Endangered Species Conservation Fund. Special nongame stamps and decals are available at Texas Parks and Wildlife Department (TPWD) field offices, most state parks, and the License Branch of TPWD headquarters in Austin. Part of the proceeds from the sale of these items is used to conserve habitat and provide information to the public concerning endangered species. Conservation organizations in Texas also welcome your participation and support.

For More Information Contact

Texas Parks and Wildlife Department
Wildlife Diversity Branch
4200 Smith School Road
Austin, Texas 78744

(512) 912-7011 or (800) 792-1112

or

U.S. Fish and Wildlife Service
Ecological Services Field Office
10711 Burnet Road, Suite 200
Austin, Texas 78758
(512) 490-0057

Management guidelines are available from the Texas Parks and Wildlife Department and U.S. Fish and Wildlife Service for landowners and managers wishing to know more about rangeland management practices which improve habitat for the Black-capped Vireo.

References

- Armstrong, W.E., M.W. Lockwood, and D.K. Stuart. 1992. *Performance report: Black-capped vireo management on Texas Parks and Wildlife Department lands*. Federal Aid Project No. E-1-4, Job No. 3.2.
- Armstrong, W.E., C. Travis, and B.G. Alexander. 1989. *Final report: Black-capped vireo management*. Federal Aid Project No. W-103-R-19, Job No. 60.
- Farquhar, C.C. and J. P. Maresh. 1998. *Population biology and habitat characterization of black-capped vireos at Dolan Falls Ranch Preserve, Val Verde County, Texas. Year two final report*. Endangered Resources Branch, Texas Parks and Wildlife Department. Austin, Texas.
- Graber, J.W. 1961. *Distribution, habitat requirements, and life history of the black-capped vireo*. *Ecol. Mon.* 31:313-336.
- Grzybowski, J.A., R.B. Clapp, and J.T. Marshall, Jr. 1986. *History and current population status of the Black-capped vireo in Oklahoma*. *American Birds* 40:1151-1161.
- Grzybowski, J.A., D.J. Tazik, and G.D. Schnell. 1994. *Regional analysis of blackcapped vireo breeding habitats*. *Condor* 96:512-544.
- Grzybowski, J.A. 1995. "Black-capped vireo (*Vireo atricapillus*).” *The birds of North America*, No. 181. The Academy of Natural Sciences, Philadelphia, and The American Ornithologist Union, Washington D.C.
- Oberholser, H.C. 1974. *The bird life of Texas*. Edgar Kincaid, Ed., University of Texas Press, Austin, Texas. Vol. 2, 1069 pp.
- Robinson, S.K., S.I. Rothstein, M.C. Brittingham, L.J. Petit, and J.A. Grzybowski. 1995. "Ecology and behavior of cowbirds and their impact on host populations" (pp 428-460) in T.E. Martin and D.M. Finch, eds. *Ecology and management of neotropical migratory birds*. Oxford University Press, New York, New York.
- U.S. Fish and Wildlife Service. 1991. *Black-capped vireo recovery plan*. Endangered Species Office, Albuquerque, N.M.
- U.S. Fish and Wildlife Service. 1996. *Black-capped vireo population and habitat viability assessment report*. Compiled and edited by Carol Beardmore, Jeff Hatfield, and Jim Lewis in conjunction with workshop participants. Report of a September 18-21, 1995 workshop arranged by the U.S. fish and Wildlife Service in partial fulfillment of U.S. National Biological Service Grant No. 80333- 1423. Austin, Texas. ix + 57 pp.

Management Guidelines for Black-capped Vireo

The following guidelines address land management practices that can be used to maintain, enhance, or create Black-capped Vireo habitat. They are intended primarily to serve as general guidance for rural landowners and others managing land for livestock and/or wildlife in Texas. The guidelines are based on our current understanding of the biology of this species.

Private landowners have a tremendous opportunity to conserve and manage the fish and wildlife resources of Texas. The objective of these guidelines is to provide landowners with recommendations about how typically-used land management practices could be conducted so that it would be unlikely that Black-capped Vireos would be impacted. The guidelines will be updated periodically to make them more practical and useful to rural landowners. The guidelines are based on the best available information and current understanding about the biology of the vireo, but may be refined as additional biological data are collected. TPWD biologists have prepared these guidelines in consultation with USFWS biologists to assure landowners who carry out land management practices within the guidelines that they would know, with the greatest certainty possible, that they would not be in violation of the Endangered Species Act.

This document also provides information on land management practices that are appropriate for protection and/or enhancement of habitat.

The categories were chosen to represent commonly encountered vegetation types and to address common questions regarding the effect of management practices on Black-capped Vireos. In addition, suggestions are offered that promote conservation of soil, water, plant, and wildlife resources.

Prescribed Burning

Fire is a natural component of Texas rangelands, and prescribed burning has many range and wildlife management benefits. These include improved forage quality and availability for livestock and deer, and maintenance of desirable plant composition and structure. Prescribed burning in some portions of the vireos range can be an excellent tool used to maintain or create the desired vegetation structure for vireo nesting; i.e. a mosaic of shrubs and open grassland with abundant woody foliage below 10 feet. If planning these activities in Bandera, Kerr, Kimble, Real, and Uvalde counties, landowners should avoid impacts to Tobusch fishhook cactus (*Ancistrocactus tobuschii*), a federally listed endangered plant, which occurs on similar soils as the vireo. Cool season burns that are patchy and low intensity, conducted prior to March 15, are often recommended to control small juniper, thus maintaining the relatively open shrublands preferred by vireos. Care should be taken to burn under appropriate humidity and wind conditions to maintain the proper black-capped vireo vegetation profile. Prescribed burns conducted during late spring and early fall, under hotter conditions, can be used to set back plant succession to create vireo habitat; however, warm season burns should be done only in areas that do not currently support Black-capped Vireos. On grazed rangeland, prescribed burns should be coordinated with livestock rotation to allow for needed deferments. It is best to avoid burning relatively small areas within large pastures to prevent heavy grazing pressure by livestock and/or deer on burned areas.

Desirable burn intervals for cool season burns vary throughout the state, depending on rainfall and vegetation type. Field experience shows that, for much of the Hill Country, a burning interval of 5 to 7 years is considered desirable to keep Ashe juniper (cedar) invasion in check and to allow regrowth of broad-leaved shrubs. Maintaining open grassland areas between clumps of shrubs is important for good vireo habitat.

Research is needed to better understand the use of prescribed burning to maintain and create vireo habitat, and to develop guidelines on desirable burn intervals throughout the vireo's range in Texas, especially in the western Edwards Plateau and eastern Trans-Pecos. Assistance from people experienced with the use of prescribed burning is highly recommended.

Landowners are encouraged to have a complete written prescribed burn plan addressing the objectives of the burn, required weather conditions, grazing deferments, fireguard preparations, personnel and equipment needed for nest concealment. Livestock and deer management, which allows woody plants such as live oak, shin oak, sumac, Texas persimmon, elbowbush, redbud, and hackberry to make dense growth from zero to at least 8 feet, is needed. On ranches throughout Texas, moderate stocking, rotation of livestock, controlling deer and exotic ungulate numbers and proper use of desirable browse plants will benefit deer and livestock as well as Black-capped Vireos.

To provide adequate nesting cover for vireos, woody plants should receive only limited browsing during the spring and summer. If animals (livestock, deer, and exotics) are wellmanaged and kept within recommended stocking rates, this can be achieved. Experience has shown that, in general, ranges stocked with cattle and deer tend to maintain better vireo nesting cover than ranges stocked with goats and exotic animals.

Limit browsing pressure, especially during the growing season, to no more than 50% of the total annual growth (current year twigs and leaves) within reach of animals on any given plant. This will maintain plants that are already vigorous and allow for improvement of those with less than ideal structure. As a rule of thumb, if you can “see through” a needed, a detailed map showing how the burn will be conducted, and notification and safety procedures.

Landowners are advised to contact local representatives of the Texas Parks and Wildlife Department, USDA Natural Resources Conservation Service, or Texas Cooperative Extension for help in developing and implementing a prescribed burning program designed specifically for your property and management objectives.

Selective Brush Management

In some portions of the vireos range, particularly the central and eastern segment, increases in juniper (cedar) and other woody species can cause the vegetation to grow out of the patchy, low shrub cover that provides suitable habitat. In these communities, good nesting habitat generally has between 30-60% shrub canopy. Selective brush removal with herbicides or mechanical means during the non-breeding season (September-February) can be used to keep the habitat favorable for vireo nesting.

For example, the selective removal of juniper, mesquite, or pricklypear (less desirable to the vireo and to the rancher) serves to maintain the proper shrub canopy and encourages growth of associated broad-leaved shrubs. Selective brush removal should strive to maintain the desired low shrubby structure. Radical changes in shrub canopy from one year to the next over large areas should be avoided, since this may alter vireo habitat too drastically within a short time-frame. However, moderate thinning of dense (>60%) shin oak so that the low canopy is maintained at 30-60% shrub canopy can enhance habitat. Western Edwards Plateau rangelands comprised primarily of mesquite, often referred to as mesquite flats, are not considered Black-capped Vireo habitat; therefore, mesquite control in these areas will not affect vireos. When using herbicides, careful attention to the kinds, amounts, timing, and application technique will achieve the best control of target species at minimum cost. Precise application also reduces the risk of environmental contamination and offsite effects. It is best to choose highly selective individual plant treatment methods, whenever practical, to avoid damage to desirable shrubs such as live oak, shin oak, Texas oak, hackberry, Texas persimmon, sumac, redbud, and elm. Herbicides should always be used in strict accordance with label directions, including those for proper storage and disposal of containers and rinse water. Herbicide applications should not occur during the breeding season, except for basal applications or

individual plant treatment of prickly pear pads.

Carefully planned mechanical methods of brush management such as chaining, roller chopping, shredding, hand cutting, hydraulic shearing, grubbing, and tree dozing can be used to achieve desirable shrub composition and to stimulate basal sprouting of key woody species in order to maintain, enhance, or create vireo habitat. If planning these activities in Bandera, Kerr, Kimble, Real, and Uvalde counties landowners should avoid impacts to Tobusch fishhook cactus (*Ancistrocactus tobuschii*), a federally listed endangered plant, which occurs on similar soils as the vireo. As with other habitat manipulation procedures, mechanical methods should only be used during the non-breeding season (September-February) and done in such a way as to maintain the proper black-capped vireo vegetation profile. Remember that good grazing management and moderate stocking rates can reduce woody plant invasion and therefore the need for expensive brush control practices.

Finally, although brush management practices can be used to change the structure and composition of vegetation so that vireos may occupy the habitat, landowners should seek technical assistance when planning brush management practices in habitat that is known to be occupied by Blackcapped Vireos. Since brush management activities can affect habitat for the Golden-cheeked Warbler as well as the Black-capped Vireo, landowners are encouraged to learn about the habitat requirements of both endangered songbirds (see TPWD leaflet on the Golden-cheeked Warbler).

Grazing and Browsing Management

Excessive browsing by goats, exotic animals, and white-tailed deer destroys the thick woody growth browse plant at “door knob” to “eye level”, then too much stem and leaf growth has been removed. Installation of structures needed to facilitate good grazing management; i.e., fencing, pipelines, water troughs, water tanks, and ponds, need to avoid removing vireo habitat, should include only enough space to allow for proper operation and maintenance, and need to conduct activities during the non-nesting period (September-February).

Careful management of woody plants will not only provide for the habitat needs of Black-capped Vireos, but will also create high quality habitat for deer and other wildlife as well as livestock. Technical assistance in identifying browse plants and determining proper use is available from the Texas Parks and Wildlife Department and USDA Natural Resources Conservation Service.

Reducing Impacts From Cowbirds

Brood parasitism by Brown-headed Cowbirds poses a serious threat to successful reproduction in some populations of Black-capped Vireos. Research is currently underway to better understand the impacts of cowbirds on vireos. Because livestock attract cowbirds, management to reduce cowbird impacts is important on grazed land.

Because cowbirds are attracted to easily available sources of food, avoid spilling or scattering grain. Supplemental feeding areas should be moved frequently and kept free from accumulations of waste grain. This would help to prevent sparsely vegetated areas of compacted soils, which also tend to attract cowbirds.

Because cowbirds can be attracted by the presence of livestock, grazing management can be used to remove grazing animals from areas where vireos nest. For example, livestock can be rotated away from prime nesting habitat during the breeding season. Another option is to graze stocker cattle during the fall and winter, resting pastures during the spring/summer nesting season. Resting pastures periodically improves range condition and may also help reduce nest

parasitism.

Finally, trapping and/or shooting cowbirds can be very effective in reducing vireo brood parasitism, since a single female cowbird can parasitize hosts over a sizeable area (4-5 acres, or more). Mounted mobile traps, placed near watering sites as livestock are rotated through pastures, have been used successfully to reduce cowbird numbers. Properly placed stationary traps have also proven effective in reducing cowbird numbers and parasitism in a local area. Shooting cowbirds at places where they congregate is another option, although this method is often not selective for the cowbirds responsible for the parasitism. Shooting female cowbirds within Black-capped Vireo nesting habitat for as little as one hour a week can reduce parasitism.

Persons trapping cowbirds need to be certified for the handling of non-target birds under the general trapping permit held by TPWD. Preventing mortality of non-target birds is very important, so traps must be carefully monitored and checked frequently.

Contact Texas Parks and Wildlife Department for information and assistance in implementing a cowbird control program.

Habitat Restoration

For landowners in central Texas wishing to restore or create habitat for the Black-capped Vireo in areas currently unoccupied by vireos, the following suggestions are offered.

One type of restorable habitat is an open shrubland capable of growing a diversity of woody plants, where much of the low-growing cover has been removed through overbrowsing by livestock or deer. Controlling browsing pressure by reducing animal numbers and providing pasture rest will allow the natural reestablishment of low-growing shrub cover needed by vireos. Prescribed burning and or mechanical methods described under the Selective Brush Management section may be needed to jump start the resprouting and root sprouting of trees and shrubs.

Habitat restoration may also be possible in areas where the shrub layer has become too tall or dense to provide good vireo habitat. In these areas, well-planned use of controlled fire or other brush management techniques listed above can reduce overall shrub height, stimulate basal sprouting of shrubs, and reduce shrub density to produce more favorable habitat for vireos. The goal is to maintain the critical low growing canopy cover of 30-60%.

Also, in areas where the brush has become too dense, selective thinning conducted during the nonnesting period (September through February) could be done to produce a more open habitat. Carefully planned brush management could be used to encourage regeneration and lateral branching of desirable shrubs by allowing sunlight to reach the ground. The idea is to restore areas to relatively open, low-growing shrub/grassland vegetation that may provide habitat preferred by vireos. If planning any of these activities in Bandera, Kerr, Kimble, Real, and Uvalde counties landowners should avoid impacts to Tobusch fishhook cactus (*Ancistrocactus tobuschii*), a federally listed endangered plant, which occurs on similar soils as the vireo.

Currently, there is no strong evidence to suggest that habitat manipulation will be necessary on many parts of the drier western and southwestern Texas range (western Edwards Plateau and eastern Trans-Pecos) as mature vegetation communities in these areas are used successfully by vireos. Unless browsing pressure or other catastrophic disturbances have eliminated desirable shrub land in these areas, the only requirement needed is time. Fire is of limited use in

lower rainfall areas devoid of fine fuels and the plant density required for cost-effective prescribed burns.

There are a number of agencies and organizations conducting management activities benefiting the vireo that can provide useful information and/or assistance to landowners. These include Texas Parks and Wildlife Department, USFWS, The Nature Conservancy, USDA Natural Resources Conservation Service, and Environmental Defense.

Summary

In the Edwards Plateau and other parts of the range supporting woodland or savanna, periodic prescribed burning and selective brush management are very effective in maintaining and creating Black-capped Vireo habitat. In all parts of the range, control of deer and exotic wildlife numbers, and good grazing management practices, including proper stocking and rotational grazing, are management options that can be used to maintain and enhance habitat for Black-capped Vireos. These same management tools will also maintain diverse and productive rangelands. In addition to providing food, fiber, and support for rural landowners, wellmanaged rangelands provide habitat for a wide variety of wildlife, and benefits such as clean water, natural diversity, and recreational opportunities for all Texans.

Technical assistance in range and wildlife management, including grazing management, determination of proper stocking rates, prescribed burning, brush management, and management for endangered species, is available to landowners and managers by contacting the Texas Parks and Wildlife Department, USDA Natural Resources Conservation Service, or Texas Cooperative Extension. Further guidance and specific questions concerning Black-capped Vireo research, endangered species management and recovery, and the Endangered Species Act, should be directed to the U.S. Fish and Wildlife Service or Texas Parks and Wildlife Department. If, after reading this leaflet, you are still unsure whether or not your management plans will adversely affect the Vireo or its habitat, please contact the U.S. Fish and Wildlife Service for assistance.