

**INTERNATIONAL ECOLOGICAL  
CLASSIFICATION STANDARD:**

**TERRESTRIAL ECOLOGICAL CLASSIFICATIONS**

**Ecological Systems of Texas' Chihuahuan Deserts**

08 October 2009

by

NatureServe

1101 Wilson Blvd., 15<sup>th</sup> floor  
Arlington, VA 22209

This subset of the International Ecological Classification Standard covers terrestrial ecological systems attributed to the Texas. This classification has been developed in consultation with many individuals and agencies and incorporates information from a variety of publications and other classifications. Comments and suggestions regarding the contents of this subset should be directed to [Judy Teague <judy\_teague@natureserve.org>].



Copyright © 2009 NatureServe, 1101 Wilson Blvd, 15<sup>th</sup> floor  
Arlington, VA 22209, U.S.A. All Rights Reserved.

**Citations:**

*The following citation should be used in any published materials which reference ecological system and/or International Vegetation Classification (IVC hierarchy) and association data:*

NatureServe. 2009. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. U.S.A. Data current as of 08 October 2009.

TABLE OF CONTENTS

**FOREST AND WOODLAND..... 4**  
 CES305.795 Madrean Encinal..... 4  
 CES305.796 Madrean Lower Montane Pine-Oak Forest and Woodland ..... 4  
 CES305.797 Madrean Pinyon-Juniper Woodland..... 4  
 CES306.813 Rocky Mountain Aspen Forest and Woodland ..... 4  
 CES306.814 Rocky Mountain Bigtooth Maple Ravine Woodland..... 5  
 CES306.835 Southern Rocky Mountain Pinyon-Juniper Woodland..... 5  
 CES306.648 Southern Rocky Mountain Ponderosa Pine Woodland ..... 5

**SHRUBLAND..... 7**  
 CES302.733 Apacherian-Chihuahuan Mesquite Upland Scrub..... 7  
 CES302.731 Chihuahuan Creosotebush Desert Scrub..... 7  
 CES302.734 Chihuahuan Mixed Desert and Thornscrub..... 7  
 CES302.017 Chihuahuan Mixed Salt Desert Scrub..... 8  
 CES302.737 Chihuahuan Stabilized Coppice Dune and Sand Flat Scrub ..... 8  
 CES302.738 Chihuahuan Succulent Desert Scrub ..... 8  
 CES303.041 Edwards Plateau Limestone Shrubland..... 9  
 CES302.031 Madrean Oriental Chaparral..... 9  
 CES306.818 Rocky Mountain Gambel Oak-Mixed Montane Shrubland ..... 9  
 CES306.822 Rocky Mountain Lower Montane-Foothill Shrubland..... 10  
 CES301.986 Tamaulipan Calcareous Thornscrub ..... 10  
 CES301.983 Tamaulipan Mixed Deciduous Thornscrub..... 10

**STEPPE/SAVANNA ..... 11**  
 CES302.735 Apacherian-Chihuahuan Semi-Desert Grassland and Steppe..... 11  
 CES302.732 Chihuahuan Gypsophilous Grassland and Steppe..... 11  
 CES301.730 Madrean Juniper Savanna ..... 11  
 CES306.649 Southern Rocky Mountain Ponderosa Pine Savanna ..... 11

**HERBACEOUS ..... 13**  
 CES302.061 Chihuahuan Loamy Plains Desert Grassland ..... 13  
 CES302.736 Chihuahuan Sandy Plains Semi-Desert Grassland ..... 13  
 CES306.824 Southern Rocky Mountain Montane-Subalpine Grassland ..... 13  
 CES303.672 Western Great Plains Shortgrass Prairie ..... 13  
 CES303.673 Western Great Plains Tallgrass Prairie ..... 14

**WOODY WETLAND ..... 15**  
 CES302.748 North American Warm Desert Lower Montane Riparian Woodland and Shrubland..... 15  
 CES302.752 North American Warm Desert Riparian Mesquite Bosque ..... 15  
 CES302.753 North American Warm Desert Riparian Woodland and Shrubland..... 15  
 CES302.755 North American Warm Desert Wash..... 15

**HERBACEOUS WETLAND ..... 16**  
 CES300.729 North American Arid West Emergent Marsh..... 16  
 CES302.747 North American Warm Desert Cienega..... 16  
 CES302.039 North American Warm Desert Interdunal Swale Wetland ..... 16  
 CES303.666 Western Great Plains Closed Depression Wetland..... 17  
 CES303.675 Western Great Plains Open Freshwater Depression Wetland..... 17  
 CES303.669 Western Great Plains Saline Depression Wetland ..... 17

**MIXED UPLAND AND WETLAND ..... 19**  
 CES302.746 Chihuahuan-Sonoran Desert Bottomland and Swale Grassland ..... 19  
 CES303.956 Western Great Plains Riparian..... 19

<b>BARREN .....</b>	<b>20</b>
<b>CES302.744 North American Warm Desert Active and Stabilized Dune .....</b>	<b>20</b>
<b>CES302.743 North American Warm Desert Badland .....</b>	<b>20</b>
<b>CES302.745 North American Warm Desert Bedrock Cliff and Outcrop.....</b>	<b>20</b>
<b>CES302.750 North American Warm Desert Pavement.....</b>	<b>20</b>
<b>CES302.751 North American Warm Desert Playa .....</b>	<b>20</b>
<b>CES302.754 North American Warm Desert Volcanic Rockland .....</b>	<b>20</b>
<b>CES306.815 Rocky Mountain Cliff, Canyon and Massive Bedrock .....</b>	<b>21</b>

## FOREST AND WOODLAND

---

### CES305.795 MADREAN ENCINAL

---

Madrean Encinal occurs on foothills, canyons, bajadas and plateaus in the Sierra Madre Occidentale and Sierra Madre Orientale in Mexico, extending north into Trans-Pecos Texas, southern New Mexico and sub-Mogollon Arizona. These woodlands are dominated by Madrean evergreen oaks along a low-slope transition below Madrean Lower Montane Pine-Oak Forest and Woodland (CES305.796) and Madrean Pinyon-Juniper Woodland (CES305.797). Lower elevation stands are typically open woodlands or savannas where they transition into desert grasslands, chaparral or in some cases deserts scrub. Common evergreen oak species include *Quercus arizonica*, *Quercus emoryi*, *Quercus intricata*, *Quercus grisea*, *Quercus oblongifolia*, *Quercus toumeyii*, and in Mexico *Quercus chihuahuensis* and *Quercus albocincta*. Madrean pine, Arizona cypress, pinyon and juniper trees may be present but do not codominate. Chaparral species such as *Arctostaphylos pungens*, *Cercocarpus montanus*, *Purshia* spp., *Garrya wrightii*, *Quercus turbinella*, *Frangula betulifolia* (= *Rhamnus betulifolia*), or *Rhus* spp. may be present but do not dominate. The graminoid layer is usually prominent between trees in grassland or steppe that is dominated by warm-season grasses such as *Aristida* spp., *Bouteloua gracilis*, *Bouteloua curtipendula*, *Bouteloua rothrockii*, *Digitaria californica*, *Eragrostis intermedia*, *Hilaria belangeri*, *Leptochloa dubia*, *Muhlenbergia* spp., *Pleuraphis jamesii*, or *Schizachyrium cirratum*, species typical of Apacherian-Chihuahuan Semi-Desert Grassland and Steppe (CES302.735). This system includes seral stands dominated by shrubby Madrean oaks typically with a strong graminoid layer. In transition areas with drier chaparral systems, stands of chaparral are not dominated by Madrean oaks; however, Madrean Encinal may extend down along drainages.

This system occurs along a low-slope transition from Madrean Pinyon-Juniper Woodland (CES305.797) or Madrean Lower Montane Pine-Oak Forest and Woodland (CES305.796).

---

### CES305.796 MADREAN LOWER MONTANE PINE-OAK FOREST AND WOODLAND

---

This system occurs on mountains and plateaus in the Sierra Madre Occidentale and Sierra Madre Orientale in Mexico, Trans-Pecos Texas, southern New Mexico and Arizona, generally south of the Mogollon Rim. These forests and woodlands are composed of Madrean pines (*Pinus arizonica*, *Pinus engelmannii*, *Pinus leiophylla*, or *Pinus strobiformis*) and evergreen oaks (*Quercus arizonica*, *Quercus emoryi*, or *Quercus grisea*) intermingled with patchy shrublands on most mid-elevation slopes (1500-2300 m elevation). Other tree species include *Cupressus arizonica*, *Juniperus deppeana*, *Pinus cembroides*, *Pinus discolor*, *Pinus ponderosa* (with Madrean pines or oaks), and *Pseudotsuga menziesii*. Subcanopy and shrub layers may include typical encinal and chaparral species such as *Agave* spp., *Arbutus arizonica*, *Arctostaphylos pringlei*, *Arctostaphylos pungens*, *Garrya wrightii*, *Nolina* spp., *Quercus hypoleucoides*, *Quercus rugosa*, and *Quercus turbinella*. Some stands have moderate cover of perennial graminoids such as *Muhlenbergia emersleyi*, *Muhlenbergia longiligula*, *Muhlenbergia virescens*, and *Schizachyrium cirratum*. Fires are frequent with perhaps more crown fires than ponderosa pine woodlands, which tend to have more frequent ground fires on gentle slopes.

---

### CES305.797 MADREAN PINYON-JUNIPER WOODLAND

---

This system occurs on foothills, mountains and plateaus in the Sierra Madre Occidentale and Sierra Madre Orientale in Mexico, Trans-Pecos Texas, southern New Mexico and Arizona, generally south of the Mogollon Rim. Substrates are variable, but soils are generally dry and rocky. The presence of *Pinus cembroides*, *Pinus discolor*, or other Madrean trees and shrubs is diagnostic of this woodland system. *Juniperus coahuilensis*, *Juniperus deppeana*, *Juniperus pinchotii*, *Juniperus monosperma*, and/or *Pinus edulis* may be present to dominant. Madrean oaks such as *Quercus arizonica*, *Quercus emoryi*, *Quercus grisea*, or *Quercus mohriana* may be codominant. *Pinus ponderosa* is absent or sparse. If present, understory layers are variable and may be dominated by shrubs or graminoids. According to USFS TES mapping (USDA 2001), *Quercus grisea* woodlands (Madrean Encinal) occur on both sides of the Guadalupe Mountains and in the southeastern portion of the Sacramento Mountains. This suggests that the associated pinyon and juniper woodlands are Madrean Pinyon-Juniper Woodland (CES305.797).

---

### CES306.813 ROCKY MOUNTAIN ASPEN FOREST AND WOODLAND

---

This widespread ecological system is more common in the southern and central Rocky Mountains but occurs in the montane and subalpine zones throughout much of the western U.S. and north into Canada. An eastern extension occurs along the Rocky Mountains foothill front and in mountain "islands" in Montana (Big Snowy and Highwood mountains), and the Black Hills of South Dakota. In California, this system is only found on the east side of the Sierra Nevada adjacent to the Great Basin. Large stands are found in the Inyo and White mountains, while small stands occur on the Modoc Plateau. Elevations generally range from 1525 to 3050 m (5000-

10,000 feet), but occurrences can be found at lower elevations in some regions. Distribution of this ecological system is primarily limited by adequate soil moisture required to meet its high evapotranspiration demand. Secondly, it is limited by the length of the growing season or low temperatures. These are upland forests and woodlands dominated by *Populus tremuloides* without a significant conifer component (<25% relative tree cover). The understory structure may be complex with multiple shrub and herbaceous layers, or simple with just an herbaceous layer. The herbaceous layer may be dense or sparse, dominated by graminoids or forbs. In California, *Symphotrichum spathulatum* (= *Aster occidentalis*) is a common forb. Associated shrub species include *Symphoricarpos* spp., *Rubus parviflorus*, *Amelanchier alnifolia*, and *Arctostaphylos uva-ursi*. Occurrences of this system originate and are maintained by stand-replacing disturbances such as avalanches, crown fire, insect outbreak, disease and windthrow, or clearcutting by man or beaver, within the matrix of conifer forests. It differs from Northwestern Great Plains Aspen Forest and Parkland (CES303.681), which is limited to plains environments. The scattered occurrences in Trans-Pecos of Texas are of interest as they represent disjunct outliers of the type occurring under highly limited circumstances.

---

#### **CES306.814 ROCKY MOUNTAIN BIGTOOTH MAPLE RAVINE WOODLAND**

---

This ecological system occurs in cool ravines, on toeslopes and slump benches associated with riparian areas in the northern and central Wasatch Range and Tavaputs Plateau extending into southern Idaho, as well as in scattered localities in southwestern Utah, central Arizona and New Mexico and the Trans-Pecos of Texas. Substrates are typically rocky colluvial or alluvial soils with favorable soil moisture. These woodlands are dominated by *Acer grandidentatum* but may include mixed stands codominated by *Quercus gambelii* or with scattered conifers. Some stands may include *Acer negundo* or *Populus tremuloides* as minor components. It also occurs on steeper, north-facing slopes at higher elevations, often adjacent to Rocky Mountain Gambel Oak-Mixed Montane Shrubland (CES306.818) or Rocky Mountain Aspen Forest and Woodland (CES306.813). In the Trans-Pecos of Texas, this system is a cool ravine woodland type generally occupying lower slope positions. It forms a patchy narrow band on lower slopes that may include *Quercus muehlenbergii*, *Acer grandidentatum*, *Fraxinus* sp., *Prunus serotina*, and *Arbutus xalapensis*.

---

#### **CES306.835 SOUTHERN ROCKY MOUNTAIN PINYON-JUNIPER WOODLAND**

---

This southern Rocky Mountain ecological system occurs on dry mountains and foothills in southern Colorado east of the Continental Divide, in mountains and plateaus of north-central New Mexico, and extends out onto limestone breaks in the southeastern Great Plains. These woodlands occur on warm, dry sites on mountain slopes, mesas, plateaus, and ridges. Severe climatic events occurring during the growing season, such as frosts and drought, are thought to limit the distribution of pinyon-juniper woodlands to relatively narrow altitudinal belts on mountainsides. Soils supporting this system vary in texture ranging from stony, cobbly, gravelly sandy loams to clay loam or clay. *Pinus edulis* and/or *Juniperus monosperma* dominate the tree canopy. *Juniperus scopulorum* may codominate or replace *Juniperus monosperma* at higher elevations. Stands with *Juniperus osteosperma* are representative the Colorado Plateau and are not included in this system. In southern transitional areas between Madrean Pinyon-Juniper Woodland (CES305.797) and Southern Rocky Mountain Pinyon-Juniper Woodland (CES306.835) in central New Mexico, *Juniperus deppeana* becomes common. Understory layers are variable and may be dominated by shrubs, graminoids, or be absent. Associated species are more typical of southern Rocky Mountains than the Colorado Plateau and include *Artemisia bigelovii*, *Cercocarpus montanus*, *Quercus gambelii*, *Achnatherum scribneri*, *Bouteloua gracilis*, *Festuca arizonica*, or *Pleuraphis jamesii*.

---

#### **CES306.648 SOUTHERN ROCKY MOUNTAIN PONDEROSA PINE WOODLAND**

---

This very widespread ecological system is most common throughout the cordillera of the Rocky Mountains, from the Greater Yellowstone region south. It is also found in the Colorado Plateau region, west into scattered locations of the Great Basin. Its easternmost extent in Wyoming is in the Bighorn Mountains. These woodlands occur at the lower treeline/ecotone between grassland or shrubland and more mesic coniferous forests typically in warm, dry, exposed sites. Elevations range from less than 1900 m in northern Wyoming to 2800 m in the New Mexico mountains. Occurrences are found on all slopes and aspects; however, moderately steep to very steep slopes or ridgetops are most common. This ecological system generally occurs on soils derived from igneous, metamorphic, and sedimentary material, with characteristic features of good aeration and drainage, coarse textures, circumneutral to slightly acidic pH, an abundance of mineral material, rockiness, and periods of drought during the growing season. Northern Rocky Mountain Ponderosa Pine Woodland and Savanna (CES306.030) in the eastern Cascades, Okanogan, and northern Rockies regions receives winter and spring rains, and thus has a greater spring "green-up" than the drier woodlands in the central Rockies. *Pinus ponderosa* (primarily var. *scopulorum* and var. *brachyptera*) is the predominant conifer; *Pseudotsuga menziesii*, *Pinus edulis*, *Pinus contorta*, *Populus tremuloides*, and *Juniperus* spp. may be present in the tree canopy. The understory is usually shrubby, with *Artemisia nova*, *Artemisia tridentata*, *Arctostaphylos patula*, *Arctostaphylos uva-ursi*, *Cercocarpus montanus*, *Purshia stansburiana*, *Purshia tridentata*, *Quercus gambelii*, *Symphoricarpos* spp., *Prunus virginiana*, *Amelanchier alnifolia* (less so in Montana), and *Rosa* spp. common species. *Pseudoroegneria spicata*, *Pascopyrum smithii*, and species of *Hesperostipa*, *Achnatherum*, *Festuca*,

*Muhlenbergia*, and *Bouteloua* are some of the common grasses. Mixed fire regimes and ground fires of variable return intervals maintain these woodlands, depending on climate, degree of soil development, and understory density.

## SHRUBLAND

---

### CES302.733 APACHERIAN-CHIHUAHUAN MESQUITE UPLAND SCRUB

---

This ecological system often occurs as invasive upland shrublands that are concentrated in the extensive desert grassland in foothills and piedmonts of the Chihuahuan Desert, extending into the Sky Island region to the west. Substrates are typically derived from alluvium, often gravelly without a well-developed argillic or calcic soil horizon that would limit infiltration and storage of winter precipitation in deeper soil layers. *Prosopis* spp. and other deep-rooted shrubs exploit this deep-soil moisture that is unavailable to grasses and cacti. Vegetation is typically dominated by *Prosopis glandulosa* or *Prosopis velutina* and succulents. Other desert scrub species that may codominate include *Acacia neovernicosa*, *Acacia constricta*, *Juniperus monosperma*, or *Juniperus coahuilensis*. *Larrea tridentata* is typically absent or has low cover. Grass cover is typically low and composed of desert grasses such as *Dasyochloa pulchella* (= *Erioneuron pulchellum*), *Muhlenbergia porteri*, *Muhlenbergia setifolia*, and *Pleuraphis mutica*. During the last century, the area occupied by this system has increased through conversion of desert grasslands as a result of drought, overgrazing by livestock, and/or decreases in fire frequency. It is similar to Chihuahuan Mixed Desert and Thorn Scrub (CES302.734) but is generally found at higher elevations where *Larrea tridentata* and other desert scrub are not codominant. It is also similar to Chihuahuan Stabilized Coppice Dune and Sand Flat Scrub (CES302.737) but does not occur on eolian-deposited substrates (sandsheets), although some stands may have evidence of wind erosion and deposition. This desert scrub occurs on substrates that are typically derived from alluvium, often gravelly without a well-developed argillic or calcic soil horizon that would limit infiltration and storage of winter precipitation in deeper soil layers. *Prosopis* spp. and other deep-rooted shrubs exploit this deep-soil moisture that is unavailable to grasses and cacti (Burgess 1995).

---

### CES302.731 CHIHUAHUAN CREOSOTEBUSH DESERT SCRUB

---

This ecological system is the common lower elevation desert scrub that occurs throughout much of the Chihuahuan Desert and has recently expanded into former desert grasslands in the northern portion of its range. Stands typically occur in flat to gently sloping desert basins and on alluvial plains, extending up into lower to mid positions of piedmont slopes (bajada). Substrates range from coarse-textured loams on gravelly plains to finer-textured silty and clayey soils in basins. Soils are alluvial, typically loamy and non-saline, and frequently calcareous as they are often derived from limestone, and to a lesser degree igneous rocks. The vegetation is characterized by a moderate to sparse shrub layer (<10% cover on extremely xeric sites) that is typically strongly dominated by *Larrea tridentata* with *Flourensia cernua* often present to codominant. A few scattered shrubs or succulents may also be present, such as *Agave lechuguilla*, *Parthenium incanum*, *Jatropha dioica*, *Koeberlinia spinosa*, *Lycium* spp., and *Yucca* spp. Additionally, *Flourensia cernua* will often strongly dominate in silty basins that are included in this ecological system. In general, shrub diversity is low as this ecological system lacks codominant thornscrub and other mixed desert scrub species that are common on the gravelly mid to upper piedmont slopes. However, shrub diversity and cover may increase locally where soils are deeper and along minor drainages with occasional *Atriplex canescens*, *Gutierrezia sarothrae*, or *Prosopis glandulosa*. Herbaceous cover is usually low and composed of grasses. Common species may include *Bouteloua eriopoda*, *Dasyochloa pulchella* (= *Erioneuron pulchellum*), *Muhlenbergia porteri*, *Pleuraphis mutica*, *Scleropogon brevifolius*, and *Sporobolus airoides*. Included in this ecological system are *Larrea tridentata*-dominated shrublands with a sparse understory that occur on gravelly to silty, upper basin floors and alluvial plains. A pebbly desert pavement may be present on the soil surface.

This ecological system is the common lower elevation desert scrub that occurs throughout much of the Chihuahuan Desert and has recently expanded into former desert grasslands in the northern portion of its range. Stands typically occur in flat to gently sloping, desert basins and on alluvial plains, extending up into the lower to mid positions of piedmont slopes (bajada). Substrates range from coarse-textured loams on gravelly plains to finer-textured silty and clayey soils in basins. Soils are alluvial, typically loamy and non-saline, and frequently calcareous as they are often derived from limestone, and to a lesser degree igneous rocks (Brown 1982, MacMahon and Wagner 1985, Henrickson and Johnston 1986, MacMahon 1988, Dick-Peddie 1993).

---

### CES302.734 CHIHUAHUAN MIXED DESERT AND THORNSCRUB

---

This ecological system is the widespread desert scrub that occurs on gravelly mid to upper bajadas, foothills and dissected gravelly alluvial fans in the Chihuahuan Desert and has recently expanded into former desert grasslands in the northern portion of its range. It generally occurs on mid to upper piedmonts above the desert plains Chihuahuan Creosotebush Desert Scrub (CES302.731) and extends up to the chaparral zone. Soils are typically well-drained, non-saline, gravelly loams often with a petrocalcic layer. Substrates are frequently derived from limestone although igneous rocks are common in some areas. Vegetation is characterized by the presence of *Larrea tridentata*, typically mixed with thornscrub or other desert scrub such as *Agave lechuguilla*, *Aloysia wrightii*, *Baccharis pteronioides*, *Dasyilirion leiophyllum*, *Flourensia cernua* (not bottomland), *Fouquieria splendens*, *Koeberlinia spinosa*, *Krameria*

*erecta*, *Leucophyllum minus*, *Mimosa aculeaticarpa* var. *biuncifera*, *Mortonia scabrella* (= *Mortonia sempervirens* ssp. *scabrella*), *Opuntia engelmannii*, *Parthenium incanum*, *Prosopis glandulosa*, and *Rhus microphylla* (in drainages). Stands of *Acacia constricta*-, *Acacia neovernicosa*- or *Acacia greggii*-dominated thornscrub are included in this system, and limestone substrates appear important for at least these species. If present, *Prosopis glandulosa* has relatively low cover and does not dominate the shrub layer. This system also includes upper piedmont stands of desert scrub that are strongly dominated by *Larrea tridentata*. Grasses are common but generally have lower cover than shrubs. Common species may include *Bouteloua curtipendula*, *Bouteloua eriopoda*, *Bouteloua gracilis*, *Bouteloua hirsuta*, *Bouteloua ramosa*, *Dasyochloa pulchella*, and *Muhlenbergia porteri*. Also included in this ecological system are shrublands with a sparse understory of *Larrea tridentata* that occur on gravelly piedmont slopes that may extend down gravelly upper basins. A pebbly desert pavement may be present on the soil surface. This may indicate remnant erosional surfaces from the early Holocene that are thought to be some of the historic distribution of *Larrea tridentata* desert scrub in the Chihuahuan Desert. Historically, much of this desert scrub was thought to be a steppe characterized by perennial desert grasses (typically *Bouteloua eriopoda*) with an open creosotebush - mixed desert shrub layer.

This ecological system is the widespread desert scrub that occurs on gravelly mid to upper bajadas, foothills and dissected gravelly alluvial fans in the Chihuahuan Desert and has recently expanded into former desert grasslands in the northern portion of its range. It generally occurs on mid to upper piedmonts above the desert plains Chihuahuan Creosotebush Desert Scrub (CES302.731) and extends up to the chaparral zone. Soils are typically well-drained, non-saline, gravelly loams often with a petrocalcic layer. Substrates are frequently derived from limestone, although igneous rocks are common in some areas (Brown 1982, MacMahon and Wagner 1985, Henrickson and Johnston 1986, MacMahon 1988, Dick-Peddie 1993).

---

### **CES302.017 CHIHUAHUAN MIXED SALT DESERT SCRUB**

---

This ecological system includes extensive open-canopied shrublands of typically saline basins in the Chihuahuan Desert. Stands often occur on alluvial flats and around playas, as well as in floodplains along the Rio Grande and Pecos rivers, possibly also extending into the San Simon of Southeastern Arizona. Substrates are generally fine-textured, saline soils. Vegetation is typically composed of one or more *Atriplex* species such as *Atriplex canescens*, *Atriplex obovata*, or *Atriplex polycarpa* along with species of *Allenrolfea*, *Flourensia*, *Salicornia*, *Suaeda*, or other halophytic plants. Graminoid species may include *Sporobolus airoides*, *Pleuraphis mutica*, or *Distichlis spicata* at varying densities.

This system includes extensive open-canopied shrublands of typically saline basins in the Chihuahuan Desert. Stands often occur on alluvial flats, around playas and floodplains of the Rio Grande and Pecos rivers, possibly also extending into the San Simon of southeastern Arizona. Sites are flat to gently sloping with slopes up to 3%. Elevation ranges from 1000-1300 m (3300-4300 feet). Substrates are generally fine-textured, saline soils but may include moderately coarse-textured alluvium in the floodplains. Water tables are generally shallow but fluctuate within reach of deep-rooted plants, and in most places are high enough that salts accumulate on the surface of the soil.

---

### **CES302.737 CHIHUAHUAN STABILIZED COPPICE DUNE AND SAND FLAT SCRUB**

---

This ecological system includes the open desert scrub of vegetated coppice dunes and sandsheets found in the Chihuahuan Desert. Stands are usually dominated by *Prosopis glandulosa* or *Artemisia filifolia* but also include *Atriplex canescens*, *Ephedra torreyana*, *Ephedra trifurca*, *Poliomintha incana*, and *Rhus microphylla* coppice and sand flat scrub usually with 10-30% total vegetation cover. *Yucca elata*, *Gutierrezia sarothrae*, *Bouteloua eriopoda*, and *Sporobolus flexuosus* are commonly present. In northern stands, *Artemisia filifolia* dominates and *Prosopis glandulosa* become less uncommon or absent. This system include degraded sandy desert plains grasslands now dominated by *Artemisia filifolia*.

---

### **CES302.738 CHIHUAHUAN SUCCULENT DESERT SCRUB**

---

This ecological system is found in the Chihuahuan Desert on colluvial slopes, upper bajadas, sideslopes, ridges, canyons, hills and mesas. Sites are hot and dry, typically with southerly aspects. Gravel and rock are often abundant on the ground surface. The vegetation is characterized by the relatively high cover of succulent species such as *Agave lechuguilla*, *Euphorbia antisyphilitica*, *Fouquieria splendens*, *Ferocactus* spp., *Opuntia engelmannii*, *Opuntia imbricata*, *Opuntia spinosior*, *Yucca baccata*, and many others. Perennial grass cover is generally low. The abundance of succulents is diagnostic of this desert scrub system, but desert shrubs are usually present. Stands in rolling topography may form a mosaic with more mesic desert scrub or desert grassland ecological systems that would occur on less xeric northerly slopes. *Agave lechuguilla* is more abundant in stands in the southern part of the mapzone. This system does not include loamy plains desert grasslands or shrub-steppe with a strong cacti component such as cholla grasslands.



---

**CES303.041 EDWARDS PLATEAU LIMESTONE SHRUBLAND**


---

This ecological system occurs as a matrix on relatively thin-soiled surfaces of plateaus of the massive limestones such as the Edwards limestone. These short to tall shrublands are variable in density depending on the relative amount of, and depth to, bedrock. *Quercus sinuata* var. *breviloba* is an important component of the system, with some areas dominated by *Quercus fusiformis*. *Juniperus ashei* is often an important component of this system. In the west, *Pinus remota* may also contribute to a scattered emergent overstory. Other shrub species may include *Rhus virens*, *Rhus lanceolata*, *Cercis canadensis* var. *texensis*, *Forestiera pubescens*, *Forestiera reticulata*, *Fraxinus texensis*, *Unghadia speciosa*, *Sophora secundiflora*, *Diospyros texana*, *Salvia ballotiflora*, *Mimosa borealis*, *Condalia hookeri*, *Rhus trilobata*, *Opuntia engelmannii*, and *Mahonia trifoliolata*. This system also includes *Quercus mohriana*- or *Quercus vaseyana*-dominated shrublands that are more common to the west, often sharing dominance with *Juniperus pinchotii*. Herbaceous cover may be patchy and is generally graminoid with species including *Schizachyrium scoparium*, *Bouteloua curtipendula*, *Bouteloua rigidisetata*, *Bouteloua trifida*, *Hilaria belangeri*, *Bothriochloa laguroides* ssp. *torreyana*, *Nassella leucotricha*, *Erioneuron pilosum*, *Aristida* spp., and others. Disturbances such as fire may be important processes maintaining this system. However, it appears to persist on thin-soiled sites. In the western portions of the Edwards Plateau, more xeric conditions lead to the slow succession of sites to woodlands, resulting in long-persisting shrublands. This system occurs on thin soils over limestone in the Edwards Plateau of Texas.

This system occurs in a steady state on thin-soiled xeric sites. Shrub cover can be 100% in patches, but overall cover may be 40-50%. Patches of dense shrubs may be interspersed with bare rock and grasslands over shallow soil. Farther west this system grades into other shallow-soiled shrubland systems.

---

**CES302.031 MADREAN ORIENTAL CHAPARRAL**


---

This ecological system occurs in mountains across southeastern New Mexico (Guadalupe Mountains), Trans-Pecos Texas (Chisos and Davis mountains) and Madrean Oriental in northern Mexico. It often dominates along the mid-elevation transition from the Chihuahuan Desert into mountains (1700-2500 m). It occurs on foothills, mountain slopes and canyons in drier habitats below the encinal and pine woodlands, and is often associated with more xeric and coarse-textured substrates such as limestone, basalt or alluvium, especially in transition areas with more mesic woodlands. The moderate to dense shrub canopy includes many shrub oak species, such as *Quercus emoryi*, *Quercus grisea*, *Quercus intricata*, *Quercus invaginata*, *Quercus laceyi*, *Quercus mohriana*, *Quercus pringlei*, *Quercus pungens*, and *Quercus vaseyana*, and several widespread chaparral species, such as *Arctostaphylos pungens*, *Ceanothus greggii*, *Cercocarpus montanus*, *Fallugia paradoxa*, and *Garrya wrightii*; other species characteristic of this system include *Arbutus xalapensis* (= *Arbutus texana*), *Fraxinus greggii*, *Fendlera rigida* (= *Fendlera linearis*), *Garrya ovata*, *Purshia mexicana*, *Rhus virens* var. *choriophylla* (= *Rhus choriophylla*), *Salvia lycioides* (= *Salvia ramosissima*), *Salvia roemeriana*, and *Salvia regla*. In the Trans-Pecos of Texas, disjunct *Quercus gambelii* may occur as a significant component of this shrubland. Most chaparral species are fire-adapted, resprouting vigorously after burning or producing fire-resistant seeds. Stands occurring within montane woodlands are seral and a result of recent fires. Grass cover may be significant. Dominant grasses often include *Bouteloua curtipendula*, *Bouteloua hirsuta*, and *Muhlenbergia emersleyi*. The similar Mogollon chaparral system has floristics mostly derived from the Sierra Madre Occidentale, whereas floristics of this system are derived from the Sierra Madre Oriental. However, this system is not matorral (thornscrub) as it is typically dominated by shrubby evergreen oaks and chaparral species, not thornscrub species. More survey is needed to determine if *Quercus turbinella*, common in the Mogollon Chaparral system, also occurs in the Madrean Oriental Chaparral.

---

**CES306.818 ROCKY MOUNTAIN GAMBEL OAK-MIXED MONTANE SHRUBLAND**


---

This ecological system occurs in the mountains, plateaus and foothills of the southern Rocky Mountains and Colorado Plateau, including the Uinta and Wasatch ranges and the Mogollon Rim. These shrublands are most commonly found along dry foothills, lower mountain slopes, and at the edge of the western Great Plains from approximately 2000 to 2900 m in elevation, and are often situated above pinyon-juniper woodlands. Substrates are variable and include soil types ranging from calcareous, heavy, fine-grained loams to sandy loams, gravelly loams, clay loams, deep alluvial sand, or coarse gravel. The vegetation is typically dominated by *Quercus gambelii* alone or codominant with *Amelanchier alnifolia*, *Amelanchier utahensis*, *Artemisia tridentata*, *Cercocarpus montanus*, *Prunus virginiana*, *Purshia stansburiana*, *Purshia tridentata*, *Robinia neomexicana*, *Symphoricarpos oreophilus*, or *Symphoricarpos rotundifolius*. There may be inclusions of other mesic montane shrublands with *Quercus gambelii* absent or as a relatively minor component. This ecological system intergrades with the lower montane-foothills shrubland system and shares many of the same site characteristics. Density and cover of *Quercus gambelii* and *Amelanchier* spp. often increase after fire.

---

**CES306.822 ROCKY MOUNTAIN LOWER MONTANE-FOOTHILL SHRUBLAND**

---

This ecological system is found in the foothills, canyon slopes and lower mountains of the Rocky Mountains and on outcrops and canyon slopes in the western Great Plains. It ranges from southern New Mexico, extending north into Wyoming, and west into the Intermountain West region. These shrublands occur between 1500 and 2900 m elevation and are usually associated with exposed sites, rocky substrates, and dry conditions, which limit tree growth. It is common where *Quercus gambelii* is absent, such as the northern Colorado Front Range and in drier foothills and prairie hills. This system is generally drier than Rocky Mountain Gambel Oak-Mixed Montane Shrubland (CES306.818) but may include mesic montane shrublands where *Quercus gambelii* does not occur. *Cercocarpus montanus* dominates pure stands in parts of Wyoming and Colorado. Scattered trees or inclusions of grassland patches or steppe may be present, but the vegetation is typically dominated by a variety of shrubs, including *Amelanchier utahensis*, *Cercocarpus montanus*, *Purshia tridentata*, *Rhus trilobata*, *Ribes cereum*, *Symphoricarpos oreophilus*, or *Yucca glauca*. Grasses are represented as species of *Muhlenbergia*, *Bouteloua*, *Hesperostipa*, and *Pseudoroegneria spicata*. Fires play an important role in this system as the dominant shrubs usually have a severe die-back, although some plants will stump sprout. *Cercocarpus montanus* requires a disturbance such as fire to reproduce, either by seed sprout or root-crown sprouting. Fire suppression may have allowed an invasion of trees into some of these shrublands, but in many cases sites are too xeric for tree growth. In Wyoming, stands where *Cercocarpus montanus* is a component of mixed shrublands are placed in Northern Rocky Mountain Montane-Foothill Deciduous Shrubland (CES306.994).

---

**CES301.986 TAMAULIPAN CALCAREOUS THORNSCRUB**

---

This xeric thornscrub ecological system is restricted to limestone and calcareous sandstone hills and caliche substrates such as along the Bordas Scarp in southern Texas and northeastern Mexico. Soils are shallow, alkaline, strongly calcareous and underlain by bedrock or a caliche layer. It has a shorter, more open shrub canopy (usually less than 2 m) when compared to more typical thornscrub growing on more favorable sites. However, shrub cover is generally greater than 70% and often greater than 85%. Dominant species include *Leucophyllum frutescens*, *Acacia berlandieri*, and *Acacia farnesiana* with many other shrub species that may be locally dominant such as *Acacia rigidula*, *Amyris madrensis*, *Amyris texana*, *Castela erecta ssp. texana*, *Celtis pallida*, *Eysenhardtia texana*, *Helietta parvifolia*, *Koeberlinia spinosa*, *Parkinsonia texana var. macra*, *Sophora secundiflora*, or *Yucca* spp. The sparse to moderately dense herbaceous layer is dominated by perennial graminoids. This system is restricted to limestone and calcareous sandstone hills and caliche substrates such as along the Bordas Scarp in southern Texas and northeastern Mexico. Soils are shallow, alkaline, strongly calcareous and underlain by bedrock or a caliche layer.

---

**CES301.983 TAMAULIPAN MIXED DECIDUOUS THORNSCRUB**

---

This thornscrub ecological system occurs throughout much of northeastern Mexico and southern Texas. It occurs on a variety of substrates and landforms. Dominant species include *Acacia roemeriana*, *Leucophyllum frutescens*, and *Prosopis glandulosa*. Other species present to codominant include *Acacia berlandieri*, *Acacia farnesiana*, *Amyris madrensis*, *Amyris texana*, *Celtis pallida*, *Parkinsonia texana*, and cacti such as *Opuntia engelmannii var. lindheimeri*.

## STEPPE/SAVANNA

---

### CES302.735 APACHERIAN-CHIHUAHUAN SEMI-DESERT GRASSLAND AND STEPPE

---

This ecological system is a broadly defined desert grassland, mixed shrub-succulent or xeromorphic oak savanna that is typical of the Borderlands of Arizona, New Mexico and northern Mexico (Apacherian region) but extends west to the Sonoran Desert, north into the Mogollon Rim and throughout much of the Chihuahuan Desert. It is found on gently sloping bajadas that support frequent fire throughout the Sky Islands and on mesas and steeper piedmont, foothill and desert mountain slopes up to 1670 m elevation in the Chihuahuan Desert. It is characterized by typically diverse perennial grasses. Common species include grasses *Bouteloua eriopoda*, *Bouteloua hirsuta*, *Bouteloua ramosa*, *Bouteloua rothrockii*, *Bouteloua curtipendula*, *Bouteloua gracilis*, *Eragrostis intermedia*, *Muhlenbergia emersleyi*, *Muhlenbergia porteri*, *Muhlenbergia setifolia*, and *Pleuraphis jamesii*, succulent species of *Agave*, *Dasyliirion*, and *Yucca*, short-shrub species of *Calliandra*, *Mimosa*, and *Parthenium*, and tall-shrub/short-tree species of *Acacia*, *Prosopis*, and various oaks (e.g., *Quercus grisea*, *Quercus emoryi*, *Quercus arizonica*, *Quercus oblongifolia*). *Pleuraphis mutica*-dominated semi-desert grasslands often with *Bouteloua eriopoda* or *Bouteloua gracilis* occurring on lowlands and loamy plains in the Chihuahuan Desert are classified as Chihuahuan Loamy Plains Desert Grassland (CES302.061). Many of the historical desert grassland and savanna areas have been converted through intensive grazing and other land uses, some to Apacherian-Chihuahuan Mesquite Upland Scrub (CES302.733) (*Prosopis* spp.-dominated). Chihuahuan grassland types that are currently included in this system are: (1) Chino grasslands of mountain slopes on acidic igneous, limestone, or deeper gravelly soils at elevations less than 1070 m (3500 feet). These sites are dominated by *Bouteloua ramosa* with *Euphorbia antisiphilitica*, *Hechtia texensis* (= *Hechtia scariosa*), *Fouquieria splendens*, *Jatropha dioica*, and *Agave lechuguilla*. (2) Desert mountain grasslands on mountain slopes between 1070 and 1370 m (3500-4500 feet) elevation on acidic igneous substrates, but also sometimes on limestone. *Bouteloua eriopoda* and *Bouteloua curtipendula* are constituents of this system. (3) Gravelly piedmont slope grasslands between 1370 and 1670 m (4500-5500 feet) elevation on Perdiz conglomerate or Tascotal tuff. These grasslands have *Bouteloua eriopoda*, *Bouteloua gracilis*, and *Dasyliirion* as common components.

---

### CES302.732 CHIHUAHUAN GYPSOPHILOUS GRASSLAND AND STEPPE

---

This ecological system is restricted to gypsum outcrops or sandy gypsiferous and/or often alkaline soils that occur in basins and slopes in the Chihuahuan Desert. Elevation range is from 1100-2000 m. These typically sparse grasslands, steppes or dwarf-shrublands are dominated by a variety of gypsophilous plants, many of which are endemic to these habitats. Characteristic species include *Tiquilia hispidissima*, *Atriplex canescens*, *Calylophus hartwegii*, *Ephedra torreyana*, *Frankenia jamesii*, *Bouteloua breviseta*, *Mentzelia perennis*, *Nama carnosum*, *Calylophus hartwegii* (= *Oenothera hartwegii*), *Selinocarpus lanceolatus*, *Sporobolus nealleyi*, *Sporobolus airoides*, and *Sartwellia flaveriae*. This system does not include the sparsely vegetated gypsum dunes that are included in North American Warm Desert Active and Stabilized Dune (CES302.744).

---

### CES301.730 MADREAN JUNIPER SAVANNA

---

This Madrean ecological system occurs in lower foothills and plains of southeastern Arizona, southern New Mexico extending into west Texas and Mexico. These savannas have widely spaced mature juniper trees and moderate to high cover of graminoids (>25% cover). The presence of Madrean *Juniperus* spp. such as *Juniperus coahuilensis*, *Juniperus pinchotii*, and/or *Juniperus deppeana* is diagnostic. *Juniperus monosperma* may be present in some stands; *Juniperus deppeana* has a broader range than this Madrean system and extends north into southern stands of Southern Rocky Mountain Juniper Woodland and Savanna (CES306.834). Stands of *Juniperus pinchotii* may be short and resemble a shrubland. Graminoid species are a mix of those found in Western Great Plains Shortgrass Prairie (CES303.672) and Apacherian-Chihuahuan Semi-Desert Grassland and Steppe (CES302.735), with *Bouteloua gracilis* and *Pleuraphis jamesii* being most common. In addition, these areas include succulents such as species of *Yucca*, *Opuntia*, and *Agave*. Juniper savanna expansion into grasslands has been documented in the last century.

---

### CES306.649 SOUTHERN ROCKY MOUNTAIN PONDEROSA PINE SAVANNA

---

This ecological system is found predominantly in the Colorado Plateau region, west into scattered locations in the Great Basin, and north along the eastern front of the southern Rocky Mountains into southeastern Wyoming. These savannas occur at the lower treeline/ecotone between grassland or shrubland and more mesic coniferous forests typically in warm, dry, exposed sites. Elevations range from less than 1900 m in central and northern Wyoming to 2800 m in the New Mexico mountains to well over 2700 m on the higher plateaus of the Southwest. It is found on rolling plains, plateaus, or dry slopes usually on more southerly aspects. This system is best described as a savanna that has widely spaced (<25% tree canopy cover) (>150 years old) *Pinus ponderosa* (primarily var.

*scopulorum* and var. *brachyptera*) as the predominant conifer. It is maintained by a fire regime of frequent, low-intensity surface fires. A healthy occurrence often consists of open and park-like stands dominated by *Pinus ponderosa*. Understory vegetation in the true savanna occurrences is predominantly fire-resistant grasses and forbs that resprout following surface fires; shrubs, understory trees and downed logs are uncommon. Important species include *Festuca arizonica*, *Muhlenbergia virescens*, *Pseudoroegneria spicata*, *Andropogon gerardii*, *Schizachyrium scoparium*, *Festuca idahoensis*, *Piptatherum micranthum*, and *Bouteloua gracilis*. A century of anthropogenic disturbance and fire suppression has resulted in a higher density of *Pinus ponderosa* trees, altering the fire regime and species composition. Presently, many stands contain understories of more shade-tolerant species, such as *Pseudotsuga menziesii* and/or *Abies* spp., as well as younger cohorts of *Pinus ponderosa*. Northern Rocky Mountain Ponderosa Pine Woodland and Savanna (CES306.030) in the eastern Cascades, Okanogan, and northern Rockies regions receives winter and spring rains, and thus has a greater spring "green-up" than the drier woodlands in the central Rockies.

## HERBACEOUS

---

### CES302.061 CHIHUAHUAN LOAMY PLAINS DESERT GRASSLAND

---

This ecological system occurs in the northern Chihuahuan Desert and extends into limited areas of the southern Great Plains on alluvial flats, loamy plains, and basins sometimes extending up into lower piedmont slopes. Sites are typically flat or gently sloping so precipitation does not run off and may be somewhat mesic if they receive runoff from adjacent areas, but these are not wetlands. Soils are non-saline, finer textured loams or clay loam. Vegetation is characterized by perennial grasses and is typically dominated by *Pleuraphis mutica* (tobosa) or with *Bouteloua eriopoda* codominant (more historically) or *Bouteloua gracilis*. In degraded stands, *Scleropogon brevifolius*, *Dasyochloa pulchella* (= *Erioneuron pulchellum*), or *Aristida* spp. may codominate. *Pleuraphis jamesii* may become important in northern stands and *Bouteloua gracilis* in the Great Plains and on degraded stands. If present, mesic graminoids such as *Pascopyrum smithii*, *Panicum obtusum*, *Sporobolus airoides*, and *Sporobolus wrightii* typically have low cover and are restricted to drainages and moist depressions (inclusions). Scattered shrubs such as *Ephedra torreyana*, *Flourensia cernua*, *Gutierrezia sarothrae*, *Larrea tridentata*, *Opuntia imbricata*, *Prosopis glandulosa*, and *Yucca* spp. may be present, especially on degraded sites.

This upland grassland is similar to the bottomland/depressional wetland system Chihuahuan-Sonoran Desert Bottomland and Swale Grassland (CES302.746) and grades into Apacherian-Chihuahuan Semi-Desert Grassland and Steppe (CES302.735) in the foothills and piedmont desert grasslands. In similar loamy plains land positions in the Great Plains, *Bouteloua gracilis*, *Buchloe dactyloides*, or *Pleuraphis jamesii* are dominant grasses in Western Great Plains Shortgrass Prairie (CES303.672). This grassland system is found from the northern to central Chihuahuan Desert and extends across the Trans-Pecos and into areas of the southwestern Great Plains. It extends from western Texas across New Mexico and into southeastern Arizona. Stands are described from Jornada del Muerto Basin, Marfa grasslands and Marathon Basin, south to central Chihuahua and Coahuila, Mexico.

Examples of this system are found on alluvial flats, loamy plains, and basins sometimes extending up into lower piedmont slopes. Sites are typically flat or gently sloping so precipitation does not run off and may be somewhat mesic if they receive runoff from adjacent areas, but these are not wetlands. Soils are non-saline, finer textured loams or clay loam.

---

### CES302.736 CHIHUAHUAN SANDY PLAINS SEMI-DESERT GRASSLAND

---

This ecological system occurs across the Chihuahuan Desert and extends into the southern Great Plains where soils have a high sand content. These dry grasslands or steppe are found on sandy plains and sandstone mesas. The graminoid layer is typically dominated or codominated by *Bouteloua eriopoda* and *Sporobolus flexuosus* with characteristic Chihuahuan species. Other common species are *Achnatherum hymenoides*, *Aristida purpurea*, *Bouteloua gracilis*, *Hesperostipa neomexicana* (minor), *Muhlenbergia arenicola*, *Pleuraphis jamesii*, *Sporobolus airoides*, *Sporobolus constrictus*, and *Sporobolus cryptandrus*. Typically, there are scattered desert shrubs and stem succulents present, such as *Ephedra torreyana*, *Ephedra trifurca*, *Opuntia imbricata*, *Yucca baccata*, *Yucca elata*, and *Yucca torreyi*, that are characteristic of the Chihuahuan Desert. The widespread shrub *Artemisia filifolia* is also frequently present, especially in the northern extent.

---

### CES306.824 SOUTHERN ROCKY MOUNTAIN MONTANE-SUBALPINE GRASSLAND

---

This Rocky Mountain ecological system typically occurs between 2200 and 3000 m elevation on flat to rolling plains and parks or on lower sideslopes that are dry, but it may extend up to 3350 m on warm aspects. Soils resemble prairie soils in that the A-horizon is dark brown, relatively high in organic matter, slightly acidic, and usually well-drained. An occurrence usually consists of a mosaic of two or three plant associations with one of the following dominant bunch grasses: *Danthonia intermedia*, *Danthonia parryi*, *Festuca idahoensis*, *Festuca arizonica*, *Festuca thurberi*, *Muhlenbergia filiculmis*, or *Pseudoroegneria spicata*. The subdominants include *Muhlenbergia montana*, *Bouteloua gracilis*, and *Poa secunda*. These large-patch grasslands are intermixed with matrix stands of spruce-fir, lodgepole pine, ponderosa pine, and aspen forests. In limited circumstances (e.g., South Park in Colorado), they form the "matrix" of high-elevation plateaus. Small-patch representations of this system do occur at high elevations of the Trans-Pecos where they present as occurrences of *Festuca arizonica* - *Blepharoneuron tricholepis* Herbaceous Vegetation (CEGL004508). These occurrences often occupy sites adjacent to Madrean Oriental Chaparral (CES302.031).

---

### CES303.672 WESTERN GREAT PLAINS SHORTGRASS PRAIRIE

---

This system is found primarily in the western half of the Western Great Plains Division in the rainshadow of the Rocky Mountains and ranges from the Nebraska Panhandle south into Texas and New Mexico, although grazing-impacted examples may reach as far north

as southern Canada where it grades into Northwestern Great Plains Mixedgrass Prairie (CES303.674). This system occurs primarily on flat to rolling uplands with loamy, ustic soils ranging from sandy to clayey. In much of its range, this system forms the matrix system with *Bouteloua gracilis* dominating this system. Associated graminoids may include *Aristida purpurea*, *Bouteloua curtipendula*, *Bouteloua hirsuta*, *Buchloe dactyloides*, *Hesperostipa comata*, *Koeleria macrantha* (= *Koeleria cristata*), *Pascopyrum smithii* (= *Agropyron smithii*), *Pleuraphis jamesii*, *Sporobolus airoides*, and *Sporobolus cryptandrus*. Although mid-height grass species may be present, especially on more mesic land positions and soils, they are secondary in importance to the sod-forming short grasses. Sandy soils have higher cover of *Hesperostipa comata*, *Sporobolus cryptandrus*, and *Yucca elata*. Scattered shrub and dwarf-dwarf species such as *Artemisia filifolia*, *Artemisia frigida*, *Artemisia tridentata*, *Atriplex canescens*, *Eriogonum effusum*, *Gutierrezia sarothrae*, and *Lycium pallidum* may also be present. Also, because this system spans a wide range, there can be some differences in the relative dominance of some species from north to south and from east to west. Large-scale processes such as climate, fire and grazing influence this system. High variation in amount and timing of annual precipitation impacts the relative cover of cool- and warm-season herbaceous species.

In contrast to other prairie systems, fire is less important, especially in the western range of this system, because the often dry and xeric climate conditions can decrease the fuel load and thus the relative fire frequency within the system. However, historically, fires that did occur were often very expansive. Currently, fire suppression and more extensive grazing in the region have likely decreased the fire frequency even more, and it is unlikely that these processes could occur at a natural scale. A large part of the range for this system (especially in the east and near rivers) has been converted to agriculture. Areas of the central and western range have been impacted by the unsuccessful attempts to develop dryland cultivation during the Dust Bowl of the 1930s. The short grasses that dominate this system are extremely drought- and grazing-tolerant. These species evolved with drought and large herbivores and, because of their stature, are relatively resistant to overgrazing. This system in combination with the associated wetland systems represents one of the richest areas for mammals and birds. Endemic bird species to the shortgrass system may constitute one of the fastest declining bird populations. In Texas, this system occurs on the Llano Estacado and ranges to but does not include the Stockton Plateau.

This system is located on primarily flat to rolling uplands. Soils typically are loamy and ustic and range from sandy to clayey. Climate is continental with mean annual precipitation generally about 300 mm ranging to 500 mm to the south in Texas. Most of the annual precipitation occurs during the growing season as thunderstorms. Precipitation events are mostly <10 cm with occasional larger events.

---

### **CES303.673 WESTERN GREAT PLAINS TALLGRASS PRAIRIE**

---

This system can be found throughout the Western Great Plains Division. It is found primarily in areas where soil characteristics allow for mesic conditions more typical of the Eastern Great Plains Division and thus are able to sustain tallgrass species. This system may be small patches interspersed within Northwestern Great Plains Mixedgrass Prairie (CES303.674) or Western Great Plains Shortgrass Prairie (CES303.672) and may also be associated with upland terraces above a floodplain system where these more mesic conditions persist. Soils are primarily loamy Mollisols that are moderately deep and rich. Those areas that contain more sandy soils should be considered part of Western Great Plains Sand Prairie (CES303.670). This system is dominated primarily by *Andropogon gerardii* and may also include *Sorghastrum nutans*, *Schizachyrium scoparium*, *Pascopyrum smithii*, *Hesperostipa spartea*, and *Sporobolus heterolepis*. *Andropogon gerardii* often dominates the lowland regions, although *Pascopyrum smithii* can be prolific if conditions are favorable. Forbs in varying density may also be present. The primary dynamics for this system include fire, climate and grazing. Fire suppression in these areas has allowed for the invasion of woody species such as *Juniperus virginiana* and *Prunus* spp. Grazing also has contributed to these changes and likewise led to a decrease of this system as overgrazing favors shortgrass and mixedgrass systems. Conversion to agriculture likewise has probably decreased the range of this system. Thus, this system likely only occurs in small patches and in scattered locations throughout the division. Large-patch occurrences are mostly isolated to slopes and swales of rolling uplands where either grazing or cultivation are more problematic. A granitic woodland association of the Wichita Mountains of Oklahoma (*Quercus fusiformis* - (*Quercus stellata*) / *Schizachyrium scoparium* Granite Woodland (CEGL004937)), formerly included here, now is included in Crosstimbers Oak Forest and Woodland (CES205.682).

This system is found primarily on loam, moderately deep, and rich Mollisols throughout the Western Great Plains Division. These soils tend to be more mesic and deep than the majority of soils within the Western Great Plains and are more typical of the Eastern Great Plains Division.

## WOODY WETLAND

---

### CES302.748 NORTH AMERICAN WARM DESERT LOWER MONTANE RIPARIAN WOODLAND AND SHRUBLAND

---

This ecological system occurs in foothill and mountain canyons and valleys of the warm desert regions of the southwestern U.S. and adjacent Mexico, and consists of mid-to low-elevation (1100-1800 m) riparian corridors along perennial and seasonally intermittent streams. Rivers include upper portions of the Gila, Santa Cruz, Salt, San Pedro, and tributaries of the lower Colorado River (below the Grand Canyon), the lower Rio Grande and Pecos (up to its confluence with Rio Hondo) that occur in the desert portions of their range. The vegetation is a mix of riparian woodlands and shrublands. Dominant trees include *Populus angustifolia*, *Populus deltoides ssp. wislizeni*, *Populus fremontii*, *Platanus wrightii*, *Juglans major*, *Fraxinus velutina*, and *Sapindus saponaria*. Shrub dominants include *Salix exigua*, *Prunus spp.*, *Alnus oblongifolia*, and *Baccharis salicifolia*. Vegetation is dependent upon annual or periodic flooding and associated sediment scour and/or annual rise in the water table for growth and reproduction.

---

### CES302.752 NORTH AMERICAN WARM DESERT RIPARIAN MESQUITE BOSQUE

---

This ecological system consists of low-elevation (<1100 m) riparian corridors along perennial and intermittent streams in valleys of the warm desert regions of the southwestern U.S. and adjacent Mexico. Rivers include the lower Colorado (within and downstream of the Grand Canyon), Gila, Santa Cruz, Salt, lower Rio Grande, Pecos (up to near its confluence with Rio Hondo), and their tributaries that occur in the desert portions of their range. Dominant trees include *Prosopis glandulosa* and *Prosopis velutina*. Shrub dominants include *Baccharis salicifolia*, *Pluchea sericea*, and *Salix exigua*. Woody vegetation is relatively dense, especially when compared to drier washes. Vegetation, especially the mesquites, tap groundwater below the streambed when surface flows stop. Vegetation is dependent upon annual rise in the water table for growth and reproduction.

---

### CES302.753 NORTH AMERICAN WARM DESERT RIPARIAN WOODLAND AND SHRUBLAND

---

This ecological system consists of low-elevation (<1200 m) riparian corridors along medium to large perennial streams throughout canyons and desert valleys of the southwestern United States and adjacent Mexico. Rivers include the lower Colorado (into the Grand Canyon), Gila, Santa Cruz, Salt, lower Rio Grande (below Elephant Butte Reservoir in New Mexico to the Coastal Plain of Texas), and the lower Pecos (up to near its confluence with Rio Hondo in southeastern New Mexico). The vegetation is a mix of riparian woodlands and shrublands. Dominant trees include *Acer negundo*, *Fraxinus velutina*, *Populus fremontii*, *Salix gooddingii*, *Salix lasiolepis*, *Celtis laevigata var. reticulata*, *Platanus racemosa*, and *Juglans major*. Shrub dominants include *Salix geyeriana*, *Shepherdia argentea*, and *Salix exigua*. Vegetation is dependent upon annual or periodic flooding and associated sediment scour and/or annual rise in the water table for growth and reproduction.

---

### CES302.755 NORTH AMERICAN WARM DESERT WASH

---

This ecological system is restricted to intermittently flooded washes or arroyos that dissect bajadas, mesas, plains and basin floors throughout the warm deserts of North America. Although often dry, the intermittent fluvial processes define this system, which are often associated with rapid sheet and gully flow. This system occurs as linear or braided strips within desert scrub-or desert grassland-dominated landscapes. The vegetation of desert washes is quite variable, ranging from sparse and patchy to moderately dense, and typically occurs along the banks, but may occur within the channel. The woody layer is typically intermittent to open and may be dominated by shrubs and small trees such as *Acacia greggii*, *Brickellia laciniata*, *Baccharis sarothroides*, *Chilopsis linearis*, *Fallugia paradoxa*, *Hymenoclea salsola*, *Hymenoclea monogyra*, *Juglans microcarpa*, *Olneya tesota*, *Parkinsonia florida*, *Prosopis spp.*, *Psoralea spinosus*, *Prunus fasciculata*, *Rhus microphylla*, *Salazaria mexicana*, or *Sarcobatus vermiculatus*. Common upland shrubs such as *Larrea tridentata* and *Ambrosia dumosa* are often present along the edges of these washes.

## HERBACEOUS WETLAND

---

### CES300.729 NORTH AMERICAN ARID WEST EMERGENT MARSH

---

This widespread ecological system occurs throughout much of the arid and semi-arid regions of western North America, typically surrounded by savanna, shrub steppe, steppe, or desert vegetation. Natural marshes may occur in depressions in the landscape (ponds, kettle ponds), as fringes around lakes, and along slow-flowing streams and rivers (such riparian marshes are also referred to as sloughs). Marshes are frequently or continually inundated, with water depths up to 2 m. Water levels may be stable, or may fluctuate 1 m or more over the course of the growing season. Water chemistry may include some alkaline or semi-alkaline situations, but the alkalinity is highly variable even within the same complex of wetlands. Marshes have distinctive soils that are typically mineral, but can also accumulate organic material. Soils have characteristics that result from long periods of anaerobic conditions in the soils (e.g., gleyed soils, high organic content, redoximorphic features). The vegetation is characterized by herbaceous plants that are adapted to saturated soil conditions. Common emergent and floating vegetation includes species of *Scirpus* and/or *Schoenoplectus*, *Typha*, *Juncus*, *Potamogeton*, *Polygonum*, *Nuphar*, and *Phalaris*. This system may also include areas of relatively deep water with floating-leaved plants (*Lemna*, *Potamogeton*, and *Brasenia*) and submergent and floating plants (*Myriophyllum*, *Ceratophyllum*, and *Elodea*).

This ecological system occurs in the arid and semi-arid regions of western North America, where semipermanently flooded habitats are found as small patches in the matrix of a relatively dry landscape. Except for stands in the semi-arid portions of the western Great Plains, emergent marsh found in the Great Plains should be classified into one of the Western Great Plains depressional wetland systems.

---

### CES302.747 NORTH AMERICAN WARM DESERT CIENEGA

---

This ecological system occurs at low elevations (<1000 m) across the warm deserts of western North America. "Cienegas" are freshwater spring-fed wetlands. The system also includes mid-elevation (1000-2000 m) wetlands found in semi-desert grasslands and Madrean evergreen woodlands. Evaporation often creates saline conditions especially on the margins as evidenced by salt-tolerant species such as *Distichlis spicata* and *Sporobolus airoides*. Typically, low-elevation examples are too warm to accumulate a deep organic layer. The type of vegetation depends on depth of water. In shallow margins, emergent plants typical of riparian vegetation are present including species of *Carex*, *Juncus*, and *Schoenoplectus*. In adjacent deeper waters, emergent marsh can be characteristic.

---

### CES302.039 NORTH AMERICAN WARM DESERT INTERDUNAL SWALE WETLAND

---

This interdunal wetland ecological system occurs in dune fields in the Chihuahuan Desert and likely in the Sonoran and Mojave deserts. This isolated or partially isolated wetland system is an occasional component of the more extensive active and stabilized desert dune system. Stands are typically small (usually less than 0.1 ha) interdunal swales that occur in wind deflation areas, where sands are scoured down to the water table. Water table may be perched over an impermeable layer of caliche or clay layer. These wetland areas are typically dominated by common emergent herbaceous vegetation, such as species of *Eleocharis*, *Juncus*, and *Schoenoplectus*, but may include endemic plants or animals. Occasionally wetlands are dominated by trees and shrubs, such as *Populus fremontii* and *Baccharis salicifolia*, which survive both being buried as dunes advance and having their root system exposed when deflation of the dune occurs. The specific dune field ecological processes distinguish these wetlands from non-dune emergent wetlands with similar species composition. Additional interdunal wetland surveys and classification work are needed at dune systems in the Chihuahuan Desert at Cuatro Cienegas, Guadalupe Mountains, and Samalayuca dunes as well as dune systems in the Sonoran or Mojave deserts, such as Algodones, Death Valley, Eureka, Gran Desierto, Kelso, Mohawk, and Salton Sea dunes, to clarify the extent of this small-patch ecological system. It may be necessary to restrict the system to the Chihuahuan Desert if that is the extent. Gypsum dunes have species unique to that substrate and may need to be treated differently.

This interdunal wetland ecological system occurs in some dune fields in the Chihuahuan and likely the Sonoran and Mojave deserts. This isolated or partially isolated wetland system is an occasional component of the more extensive active and stabilized desert dune system. Stands are typically small (usually less than 0.1 ha) interdunal swales that occur in wind deflation areas, where sands are scoured down to the water table. Water table may be perched over an impermeable layer of caliche or clay. Dune sands may be quartz or gypsum. Gypsum dunes have species unique to that substrate and may need to be treated differently. The specific dune field ecological processes distinguish these wetlands from similar emergent wetlands. In west Texas, these wetlands occur in interdunal swales in Monahan and Kermit quartz sand dunes. These dunes occur northeast of the Pecos River where the prevailing southwest winds have blown the sands to the east where they are trapped by the escarpment of the High Plains (Southern Shortgrass Prairie Ecoregion). The ponds are on a perched water table underlain by impermeable caliche layers. The White Sands dunes of the Tularosa



Basin in southern New Mexico are a gypsum interdunal/dune system that is moving/semi-stable; during the rainy season, many of the interdunes become ephemeral lakes with wetland indicators.

---

### **CES303.666 WESTERN GREAT PLAINS CLOSED DEPRESSION WETLAND**

---

Communities associated with the playa lakes in the southern areas of this province and the rainwater basins in Nebraska characterize this system. They are primarily upland depressional basins. This hydric system is typified by the presence of an impermeable layer such as a dense clay, hydric soil and is usually recharged by rainwater and nearby runoff. They are rarely linked to outside groundwater sources and do not have an extensive watershed. Ponds and lakes associated with this system can experience periodic drawdowns during drier seasons and years, and are often replenished by spring rains. *Eleocharis* spp., *Hordeum jubatum*, along with common forbs such as *Coreopsis tinctoria*, *Symphotrichum subulatum* (= *Aster subulatus*), and *Polygonum pensylvanicum* (= *Polygonum bicorne*) are common vegetation in the wetter and deeper depression, while *Pascopyrum smithii* and *Buchloe dactyloides* are more common in shallow depressions in rangeland. Species richness can vary considerably among individual examples of this system and is especially influenced by adjacent land use, which is often agriculture, and may provide nutrient and herbicide runoff. Dynamic processes that affect these depressions are hydrological changes, grazing, and conversion to agricultural use. Open and emergent marshes may be a separate system from wet meadows and wet prairies. This system needs to be more clearly distinguished from the similar open depressional wetlands of the western Great Plains, as well as from Great Plains Prairie Pothole (CES303.661).

This system is typified by upland depressional basins with an impermeable layer such as dense clay, hydric soils. Rainwater and runoff primarily recharge this system and it is rarely linked to outside groundwater sources.

---

### **CES303.675 WESTERN GREAT PLAINS OPEN FRESHWATER DEPRESSION WETLAND**

---

This Great Plains emergent marsh ecological system is composed of lowland depressions; it also occurs along lake borders that have more open basins and a permanent water source through most of the year, except during exceptional drought years. These areas are distinct from Western Great Plains Closed Depression Wetland (CES303.666) by having a large watershed and/or significant connection to the groundwater table. A variety of species are part of this system, including emergent species of *Typha*, *Carex*, *Eleocharis*, *Juncus*, *Spartina*, and *Schoenoplectus*, as well as floating genera such as *Potamogeton*, *Sagittaria*, *Stuckenia*, or *Ceratophyllum*. The system includes submergent and emergent marshes and associated wet meadows and wet prairies. These types can also drift into stream margins that are more permanently wet and linked directly to the basin via groundwater flow from/into the pond or lake. Some of the specific communities will also be found in the floodplain system and should not be considered a separate system in that case. These types should also not be considered a separate system if they are occurring in lowland areas of the prairie matrix only because of an exceptional wet year.

This system occurs widely throughout the western Great Plains, but in the arid shortgrass region, it is replaced by North American Arid West Emergent Marsh (CES300.729). Open and emergent marshes may be a separate system from wet meadows and wet prairies. More clarification needs to be made between this system and other depressional wetlands occurring in Wyoming and Montana, such as the Inter-Mountain Basins Alkaline Closed Depression (CES304.998), Great Plains Prairie Pothole (CES303.661), and the other western Great Plains depressional wetland systems.

This system is found within lowland depressions and along lakes that have more permanent water sources throughout the year. These areas typically have a large watershed and are connected to the groundwater sources. Examples may also drift into stream margins that are more permanently wet and linked to a basin via groundwater flow from/into a pond or lake. Those areas that are found within larger prairie matrix that are only lowland or wet because of an exceptional wet year are not part of this system.

---

### **CES303.669 WESTERN GREAT PLAINS SALINE DEPRESSION WETLAND**

---

This ecological system is very similar to Western Great Plains Open Freshwater Depression Wetland (CES303.675) and Western Great Plains Closed Depression Wetland (CES303.666). However, strongly saline soils cause both the shallow lakes and depressions and the surrounding areas to be more brackish. Salt encrustations can occur on the surface in some examples of this system, and the soils are severely affected and have poor structure. Species that typify this system are salt-tolerant and halophytic species such as *Distichlis spicata*, *Sporobolus airoides*, and *Hordeum jubatum*. Other commonly occurring taxa include *Puccinellia nuttalliana*, *Salicornia rubra*, *Schoenoplectus maritimus*, *Schoenoplectus americanus*, *Suaeda calceoliformis*, *Spartina* spp., *Triglochin maritima*, and shrubs such as *Sarcobatus vermiculatus* and *Krascheninnikovia lanata*. During exceptionally wet years, an increase in precipitation can dilute the salt concentration in the soils of some examples of this system which may allow for less salt-tolerant species to occur. Communities found within this system may also occur in floodplains (i.e., more open depressions) but probably should not be considered a separate system unless they transition to areas outside the immediate floodplain. Open and emergent saline marshes may be a separate system from saline wet meadows and prairies. This system is often intimately associated (in space) with

greasewood flats, and there is some overlap in the associations between the two. This system tends to be more of an herbaceous wetlands, whereas Inter-Mountain Basins Greasewood Flat (CES304.780) is more strongly shrub-dominated with patches of herb-dominance.

This system is distinct from the freshwater depression systems by its brackish nature caused by strongly saline soils. Salt encrustations could occur near the surface in some examples of this system.

## MIXED UPLAND AND WETLAND

---

### CES302.746 CHIHUAHUAN-SONORAN DESERT BOTTOMLAND AND SWALE GRASSLAND

---

This ecological system occurs in relatively small depressions or swales and along drainages throughout the northern and central Chihuahuan Desert and adjacent Sky Islands and Sonoran Desert, as well as limited areas of the southern Great Plains on broad mesas, plains and valley bottoms that receive runoff from adjacent areas. Occupying low topographic positions, these sites generally have deep, fine-textured soils that are neutral to slightly or moderately saline/alkaline. During summer rainfall events, ponding is common. Vegetation is typically dominated by *Sporobolus airoides*, *Sporobolus wrightii*, *Pleuraphis mutica* (tobosa swales), or other mesic graminoids such as *Pascopyrum smithii* or *Panicum obtusum*. With tobosa swales, sand-adapted species such as *Yucca elata* may grow at the swale's edge in the deep sandy alluvium that is deposited there from upland slopes. *Sporobolus airoides* and *Sporobolus wrightii* are more common in alkaline soils and along drainages. Other grass species may be present, but these mesic species are diagnostic. Scattered shrubs such as *Atriplex canescens*, *Prosopis glandulosa*, *Ericameria nauseosa*, *Fallugia paradoxa*, *Krascheninnikovia lanata*, or *Rhus microphylla* may be present. This bottomland/depressional wetland system can be similar to the upland Chihuahuan Loamy Plains Desert Grassland (CES302.061) but is restricted to moist depressions and intermittently flooded drainage terraces and adjacent flats. Alkali sacaton (*Sporobolus airoides*) is often associated with more alkaline (to gypsic), poorly drained areas and giant sacaton (*Sporobolus wrightii*) with less alkaline better drained areas. *Distichlis spicata*, *Allenrolfea occidentalis*, and *Suaeda* spp. are characteristic of more saline and alkaline sites.

This ecological system occurs in relatively small depressions or swales and along drainages on broad mesas, plains and valley bottoms that receive runoff from adjacent areas. These sites occupy low topographic positions and generally have deep, fine-textured soils that are neutral to slightly or moderately saline/alkaline.

---

### CES303.956 WESTERN GREAT PLAINS RIPARIAN

---

This ecological system is found in the riparian areas of medium and small rivers and streams throughout the western Great Plains. It is likely most common in the Shortgrass Prairie and Northern Great Plains Steppe but extends west as far as the Rio Grande in New Mexico and into the Wyoming Basins in the north. Major rivers include the North and South Platte, portions of the Arkansas, Cimarron, Canadian and upper Pecos rivers and tributaries to where they extend into Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland (CES306.821). It is found on alluvial soils in highly variable landscape settings, from deep cut ravines to wide, braided streambeds. Hydrologically, these sites tend to be more flashy with less developed floodplains than on larger rivers that are classified as floodplain systems, and may dry down completely for some portion of the year. Water sources for this riparian system are largely snowmelt near the Rocky Mountains, but it will respond to summer rains. This system includes numerous smaller prairie rivers and streams that are often groundwater-fed, such as the Arikaree River, a tributary of the Republican River. Dominant vegetation shares much with generally drier portions of larger floodplain systems downstream, but overall abundance of vegetation is generally lower. Communities within this system range from riparian forests and shrublands to gravel/sand flats. Dominant species include *Populus deltoides*, *Salix* spp., *Artemisia cana* ssp. *cana*, *Pascopyrum smithii*, *Panicum virgatum*, *Panicum obtusum*, *Sporobolus cryptandrus*, and *Schizachyrium scoparium*. On the North Platte in southeastern Wyoming, *Fraxinus pennsylvanica* may be present to dominant. These areas are often subjected to heavy grazing and/or agriculture and can be heavily degraded. *Tamarix* spp., *Elaeagnus angustifolia*, and less desirable grasses and forbs can invade degraded examples up through central Colorado. Groundwater depletion and lack of fire have resulted in additional species changes. All the riparian/floodplain/alluvial systems of the Great Plains region need to be revisited for naming conventions, along with better definitions of conceptual boundaries. There is much apparent overlap in their concepts and distribution, and the names add to the confusion. In particular, the difference between "riparian" and "floodplain" usage in the names needs revisiting and possible changing. These systems include Northwestern Great Plains Floodplain (CES303.676), Northwestern Great Plains Riparian (CES303.677), Western Great Plains Floodplain (CES303.678), and Western Great Plains Riparian (CES303.956).

## BARREN

---

### CES302.744 NORTH AMERICAN WARM DESERT ACTIVE AND STABILIZED DUNE

---

This ecological system occurs across the warm deserts of North America and is composed of unvegetated to sparsely vegetated (generally <10% plant cover) active dunes and sandsheets derived from quartz or gypsum sands. Common vegetation includes *Ambrosia dumosa*, *Abronia villosa*, *Artemisia filifolia*, *Atriplex canescens*, *Eriogonum deserticola*, *Larrea tridentata*, *Pleuraphis rigida*, *Poliomntha* spp., *Prosopis* spp., *Psorothamnus* spp., *Rhus microphylla*, and *Sporobolus flexuosus*. Dune "blowouts" and subsequent stabilization through succession are characteristic processes.

---

### CES302.743 NORTH AMERICAN WARM DESERT BADLAND

---

This ecological system is restricted to barren and sparsely vegetated (generally <10% plant cover) substrates typically derived from marine shale or mudstone (badlands and mudhills). The harsh soil properties and high rate of erosion and deposition are driving environmental variables supporting sparse shrubs and dwarf-shrubs e.g., *Atriplex hymenelytra*, and herbaceous vegetation.

---

### CES302.745 NORTH AMERICAN WARM DESERT BEDROCK CLIFF AND OUTCROP

---

This ecological system is found from subalpine to foothill elevations and includes barren and sparsely vegetated landscapes (generally <10% plant cover) of steep cliff faces, narrow canyons, and smaller rock outcrops of various igneous, sedimentary, and metamorphic bedrock types. Also included are unstable scree and talus slopes that typically occur below cliff faces. Species present are diverse and may include *Bursera microphylla*, *Fouquieria splendens*, *Nolina bigelovii*, *Opuntia bigelovii*, and other desert species, especially succulents. Lichens are predominant lifeforms in some areas. May include a variety of desert shrublands less than 2 ha (5 acres) in size from adjacent areas.

---

### CES302.750 NORTH AMERICAN WARM DESERT PAVEMENT

---

This ecological system occurs throughout much of the warm deserts of North America and is composed of unvegetated to very sparsely vegetated (<2% plant cover) landscapes, typically flat basins where extreme temperature and wind develop ground surfaces of fine to medium gravel coated with "desert varnish." This sparsely vegetated system may surround playas in valley bottoms or near washes and, less commonly, on dissected, eroding alluvial fans. Very low cover of desert scrub species such as *Larrea tridentata* or *Eriogonum fasciculatum* is usually present. However, ephemeral herbaceous species may have high cover in response to seasonal precipitation, including *Chorizanthe rigida*, *Eriogonum inflatum*, and *Geraea canescens*.

---

### CES302.751 NORTH AMERICAN WARM DESERT PLAYA

---

This ecological system is composed of barren and sparsely vegetated playas (generally <10% plant cover) found across the warm deserts of North America, extending into the extreme southern end of the San Joaquin Valley in California. Playas form with intermittent flooding, followed by evaporation, leaving behind a saline residue. Salt crusts are common throughout, with small saltgrass beds in depressions and sparse shrubs around the margins. Subsoils often include an impermeable layer of clay or caliche. Large desert playas tend to be defined by vegetation rings formed in response to salinity. Given their common location in windswept desert basins, dune fields often form downwind of large playas. In turn, playas associated with dunes often have a deeper water supply. Species may include *Allenrolfea occidentalis*, *Suaeda* spp., *Distichlis spicata*, *Eleocharis palustris*, *Oryzopsis* spp., *Sporobolus* spp., *Tiquilia* spp., or *Atriplex* spp. Ephemeral herbaceous species may have high cover periodically. Adjacent vegetation is typically Sonora-Mojave Mixed Salt Desert Scrub (CES302.749), Chihuahuan Mixed Salt Desert Scrub (CES302.017), Gulf of California Coastal Mixed Salt Desert Scrub (CES302.015), Baja California del Norte Gulf Coast Ocotillo-Limberbush-Creosotebush Desert Scrub (CES302.014), or Chihuahuan Creosotebush Desert Scrub (CES302.731).

---

### CES302.754 NORTH AMERICAN WARM DESERT VOLCANIC ROCKLAND

---

This ecological system occurs across the warm deserts of North America and is restricted to barren and sparsely vegetated (<10% plant cover) volcanic substrates such as basalt lava (malpais) and tuff. Vegetation is variable and includes a variety of species depending on local environmental conditions, e.g., elevation, age and type of substrate. Typically scattered *Larrea tridentata*, *Atriplex hymenelytra*, or other desert shrubs are present.

---

**CES306.815 ROCKY MOUNTAIN CLIFF, CANYON AND MASSIVE BEDROCK**

---

This ecological system of barren and sparsely vegetated landscapes (generally <10% plant cover) is found from foothill to subalpine elevations on steep cliff faces, narrow canyons, and smaller rock outcrops of various igneous (intrusives), sedimentary, and metamorphic bedrock types. It is located throughout the Rocky Mountains and northeastern Cascade Ranges in North America. Also included are unstable scree and talus slopes that typically occur below cliff faces. In general these are the dry sparsely vegetated places on a landscape. The biota on them reflect what is surrounding them, unless it is an extreme parent material. There may be small patches of dense vegetation, but it typically includes scattered trees and/or shrubs. Characteristic trees include species from the surrounding landscape, such as *Pseudotsuga menziesii*, *Pinus ponderosa*, *Pinus flexilis*, *Populus tremuloides*, *Abies concolor*, *Abies lasiocarpa*, or *Pinus edulis* and *Juniperus* spp. at lower elevations. There may be scattered shrubs present, such as species of *Holodiscus*, *Ribes*, *Physocarpus*, *Rosa*, *Juniperus*, and *Jamesia americana*, *Mahonia repens*, *Rhus trilobata*, or *Amelanchier alnifolia*. Soil development is limited, as is herbaceous cover. This has a very broad elevation range (<3350 m) for a system; consider dividing into foothills/montane and subalpine. And/or by floristic division. This is in the Okanagan and Rockies as the montane sparse. North Pacific Montane Massive Bedrock, Cliff and Talus (CES204.093) includes everything in the Cascades and west, except the northeastern Cascades, where occurrences are this system (CES306.815). Inter-Mountain Basins Cliff and Canyon (CES304.779) occurs in the dry foothills on the east side of EDC MapZone1.

This system is located throughout the Rocky Mountain, including the isolated island ranges of central Montana, and northeastern Cascade Ranges in North America.