

## APPENDIX T

# Nongame Wildlife Management Recommendations

Following is a list and brief description of management practices that are beneficial to nongame species of wildlife. It should be noted that many of the practices are also beneficial to and recommended for game species (eg. deer, dove, turkey quail, etc.). Conversely, most management practices directed at managing game species will also be beneficial to many species of nongame wildlife.

### HABITAT CONTROL

**Prairie/grassland restoration** - Establishing a mixture of native grasses and forbs on disturbed range or farm land to provide habitat for wildlife diversity. Use the TPWD wildscape plant list.

**Forest/woodland restoration** - Establishing native trees and shrubs where appropriate to restore native habitats for wildlife diversity. Use the TPWD wildscape plant list.

**Shrubland restoration** - Establishing native shrubs or small trees where appropriate to restore native habitats for wildlife diversity. Use TPWD the wildscape plant list.

**Wetland restoration** - Establishing water flows and native vegetation in altered coastal and inland wetlands to provide wildlife habitat.

**Riparian area management** - Provide alternate livestock feeding and watering sites, exclude pastures with riparian areas from livestock grazing or fence out livestock. Defer grazing in riparian areas during April - October.

**Prescribed burning** - The use of fire to restore, enhance or maintain native habitats for wildlife diversity.

**Mowing** - Used to manage invading woody plants and maintain desirable herbaceous vegetation for wildlife food and cover. Mow before or after nesting season to avoid grassland nesting birds (most nesting occurs generally April-June).

**Exotic or "weedy" plant control** - Use of fire, selective herbicides, and mechanical methods to control invasive plants in important habitat types to maintain or restore wildlife populations.

**Conversion of exotic vegetation** - Removal and replacement of exotic vegetation with native plants for wildlife habitat.

**Restore and maintain native prairie/grassland** - Prescribed burns should be conducted according to TPWD, USDA Natural Resource Conservation Service, Texas Agricultural Extension Service and Texas Natural Resource Conservation Commission

protocols in coordination with local Fire Department. Most burns are conducted during December-March. Late winter-early spring burns will not impact cool season forbs as much as mid-winter burns. Summer burns are more risky, but could be more effective at woody plant control. If mechanical brush control is used leave brush piles for small mammals. Reseed areas with native grass/forb mixtures as necessary.

**Enhance mid-succession brush habitat** - Promote brush regeneration with prescribed fire and/or mechanical methods that remove the top-growth of woody plants but encourage root sprouting. Use proper grazing management.

**Protect karst, caves and other underground resources** - Construct appropriate cave gates or other features to minimize human disturbance to roosting bats. Insure quality underground water resources through proper disposal of toxicants and runoff management. Maintain unobstructed cave entrance for easy access by bats.

## **EROSION CONTROL**

**Riparian area management** - Provide alternate livestock feeding and watering sites, exclude pastures with riparian areas from livestock grazing or fence out livestock. Defer grazing in riparian areas during April - October. Control erosion using water structures and native plants.

**Prairie/grassland restoration** - Establishing a mixture of native grasses and forbs on disturbed range or farm land to provide habitat for wildlife diversity. Use the TPWD wildscape plant list.

**Forest/woodland restoration** - Establishing native trees and shrubs where appropriate to restore native habitats for wildlife diversity. Use the TPWD wildscape plant list.

**Trails and signs** - Create walkways or paths to manage human impact and reduce erosion in sensitive areas.

## **PREDATOR CONTROL**

**Avian predator and parasite control** - Reduce the impact of selected avian predators (grackles, ravens, crows) and brown-headed cowbirds on nesting birds through shooting and trapping, grazing management, and maintenance of large blocks of wildlife habitat.

**Carnivore-furbearer control** - Reduce the impact of coyotes, raccoons and other carnivores on colonial nesting birds.

## **PROVIDING SUPPLEMENTAL WATER**

**Wetland restoration** - Establishing moist soil management in playa wetlands (High Plains) and managing wet meadow/perched water table/riparian habitats (Rolling

Plains).

**Well/trough/pond with overflows** - Establish additional shallow water supplies through construction of ground-level wildlife ponds, or adding overflow systems on existing wells and troughs. Protect these areas from livestock use.

### **PROVIDING SUPPLEMENTAL FOOD**

**Prairie/grassland restoration** - Establishing a mixture of native grasses and forbs on disturbed range or farm land to provide habitat for wildlife diversity. Use the TPWD wildscape plant list.

**Woody restoration** - Establishing native trees and shrubs where appropriate to restore native habitats for wildlife diversity. Use the TPWD wildscape plant list.

**Butterfly and hummingbird gardens** - Establish native wildflowers, trees, shrubs, vines, or cultivated flowers as food sources for butterflies and hummingbirds. Use the TPWD wildscape plant list.

**Feeding stations** - Set up liquid, seed and free-choice feeding stations for resident and migratory birds. Especially critical during migration and winter months when natural food sources are scarce.

**Reduction of broadcast insecticides** - Increases the amount of insects available as a wildlife food source for birds, reptiles and amphibians.

**Conversion of exotic vegetation** - Removal and replacement of exotic vegetation with native plants for wildlife habitat.

### **PROVIDING SUPPLEMENTAL SHELTER**

**Brush piles/rock piles** - Leaving or stacking cleared brush and rock to create denning and escape cover for birds, small mammals, reptiles and amphibians.

**Thickets of native brush** - Create or maintain thickets of native shrubs/trees for refuge.

**Prairie/grassland restoration** - Establishing a mixture of native grasses and forbs on disturbed range or farm land to provide habitat for wildlife diversity. Use the TPWD wildscape plant list.

**Snag maintenance and creation** - Protect snags and deadfall for cavity dwelling species. Create snags using selective herbicides or girdling undesirable woody plants.

**Nest boxes and perching platforms/poles** - Provide nest structures for songbirds, owls, small mammals, bats, raptors, herons, and other nongame species.

## **CENSUS (Surveys)**

**Time area counts** - The number of individual species seen or heard during a fixed time frame per unit area (eg. point counts for birds, squirrels).

**Drift fences/pit fall traps** - A system of flashing or similar material arranged on the ground to funnel small wildlife species into buried buckets or other pitfall trap. (used primarily for reptiles and amphibians).

**Small mammal traps** - Small live traps arranged along a trapline to sample small mammals.

**Other or Indicator Species:** Bobwhite quail, dove, and wild turkey may be desired game species to have in the area, which may be expressed in the overall objective. The land management techniques that have been recommended primarily for the deer population can benefit these game birds and many other wildlife species also. These are: prescribed burning, disking, cattle rotation or exclusion from woods and certain native grass areas during certain periods, and supplemental food plots. See Appendix G for more information on quail and Appendix I for wild turkey.

**Nest/Roost boxes for Cavity Nesters/Roosters:** Where suitable nest cavities are in short supply due to lack of dead timber snags that provide cavities or natural timber hollows, artificial nest/roost boxes can be erected to help alleviate these shortages for particular species. Some of the birds and mammals that can benefit from these structures are: bluebirds, chickadees, titmice, prothonotary warbler, wrens, woodpeckers, screech owls, kestrels, wood ducks, black-bellied whistling ducks, squirrels, and bats. The TPWD Nongame and Urban Program can furnish additional information regarding number, specifications, placement, and maintenance of these structures for specific species.

**Neotropical Migratory Birds:** These are birds that breed in the United States and Canada, and migrate to the Neotropical regions of Mexico, Central and South America, and the Caribbean during the nonbreeding season. As mentioned in the General Habitat Management section at the beginning of this example plan, loss and fragmentation of woodland and native grassland habitat has reduced populations of many neotropical populations. Neotropicals include the following groups of birds: kites, hawks, falcons, owls, cuckoos, nightjars, hummingbirds, flycatchers, swallows, thrushes, vireos, warblers, tanagers, grosbeaks, buntings, sparrows, orioles, and blackbirds. For more information regarding neotropical status, surveys, and possible management strategies, contact the Partners in Flight Program Coordinator at TPWD Headquarters in Austin.

**Waterfowl/Wading Birds:** To improve the habitat for dabbling ducks and wading birds, construction of 3 - 4 foot high levees with a drop-board water control structure in suitable low areas could back up and hold water during the fall, winter, spring, summer months, depending on water management strategy. This could provide shallow (6 to 24 inches) water feeding areas for migrant ducks, wading birds, and spring-nesting wood

ducks. Exclude livestock from this area with installation of an electric or barbed wire fence around the perimeter, at least 50 yards away from the maximum flooded area. Contact the local Natural Resources Conservation Service or TPWD waterfowl biologist for assistance in location and construction of the levee.

Installation of wood duck nest boxes in and around the edge of shallow water areas can increase nesting sites for wood ducks that are normally present in the summer, but lack suitable nesting sites due to lack of natural cavities in older, damaged trees or lack of these types of trees. One nest box (not within view of other nest boxes) per acre of brood-rearing wetland habitat is usually sufficient. These should be erected on 10 foot metal or treated wooden posts in or at the edge of wetlands.

**Feral Hogs** should be controlled by shooting and live trapping whenever possible. Most success at this usually occurs during the winter when feral hogs are having to travel more to find food. Besides rooting up pastures, feral hogs compete directly with deer, turkey and most other wildlife species that rely heavily on acorns and other hard and soft mast for winter food. Deer also tend to avoid areas when feral hogs are present.

**Other Comments:** The development of a Landowner Wildlife Management Association with adjacent and neighboring landowners will greatly enhance any management that you apply to your ranch, and is strongly encouraged. TPWD and TCE personnel are available to assist in this endeavor.