

# Growth and Survival of Cage-Held Eastern Oysters in Corpus Christi Bay


by

Joseph D. Gray      Timothy L. King  
William B. Kehoe    Robert L. Colura  
Gary C. Matlock

Management Data Series  
No. 48  
1990



TEXAS  
PARKS & WILDLIFE  
DEPARTMENT  
FISHERIES DIVISION  
4200 Smith School Road  
Austin, Texas 78744



GROWTH AND SURVIVAL OF CAGE-HELD EASTERN  
OYSTERS IN CORPUS CHRISTI BAY

by

Joseph D. Gray, Timothy L. King, William B. Kehoe  
Robert L. Colura, and Gary C. Matlock

MANAGEMENT DATA SERIES

No. 48

1990

Texas Parks and Wildlife Department  
Fisheries Division  
Coastal Fisheries Branch  
4200 Smith School Road  
Austin, Texas 78744

## ACKNOWLEDGEMENTS

The authors would like to thank Phil Hale and the 4600 Building Homeowners Association for the use of their facilities as a study location. We would also like to thank Tom Heffernan, Larry McEachron, Joe Martin and Billy Fuls for their assistance in determining the study site, Paul Hammerschmidt for his review of the manuscript, and Dr. Sammy Ray of Texas A&M University for analyzing the oysters for Perkinsus marinus. We would also like to thank Rose Cortez for typing the manuscript.

## ABSTRACT

The Eastern oyster (Crassostrea virginica) was commercially harvested from Corpus Christi Bay until 1959, after which most reefs were non-productive. Reasons for the demise of oyster populations in Corpus Christi Bay are unknown, but may be related to increased salinities. Re-establishment of an Eastern oyster population in Corpus Christi Bay may be possible through stocking of oysters from other bays. The purpose of this study was to compare growth and survival of Eastern oysters from six Texas bay systems held in Corpus Christi Bay. Eastern oysters from Galveston, Matagorda, San Antonio, Aransas, Nueces (Corpus Christi Bay system), and South Bays were maintained in cages in Corpus Christi Bay for 153 days. Final results suggest no statistically significant differences in survival or mean total length among bay systems. South Bay oysters exhibited significantly higher growth rate than San Antonio and Aransas bay oysters at 56 and 153 days respectively. Results suggest no distinct advantage in stocking Eastern oysters from bays other than the Corpus Christi Bay system.



## INTRODUCTION

Oysters have occupied their current niche for some 350 million years while undergoing little evolutionary change (Longwell and Stiles 1970). Fossil evidence suggests oysters were one of the most predominant inshore inhabitants of the Atlantic coasts of North America and Europe (Yonge 1960). However, overharvesting, pollution, poorly managed cultivation practices, general habitat loss and other anthropogenic factors have contributed to the oyster's decline (Longwell and Stiles 1970).

The Eastern oyster (Crassostrea virginica) is commercially harvested from public reefs in most bay systems along the Texas gulf coast. Galveston, Matagorda and San Antonio Bays contain about 93% of the public reefs. No commercially harvestable reefs exist south of Nueces Bay except in South Bay, and these reefs constitute < 1% of the commercial landings (Quast et al. 1988).

Corpus Christi Bay once supported a commercial Eastern oyster fishery. However, commercial landings steadily declined until most reefs were non-productive by 1959 (Martinez 1963, 1964). Reasons for the demise of the populations in Corpus Christi Bay are unknown. Persistent salinity levels exceeding 25 o/oo (attributed to reduced freshwater inflow), inadequate substrate, and overfishing may have been contributing factors.

Current management practices used to restore depleted oyster fisheries range from harvest restrictions and artificial reef construction strategies to re-seeding programs which supplement local populations. Additionally, transplanting adults for spawning purposes has been used to enhance larval recruitment (Malinowski and Whitlatch 1988). Mortality studies in Aransas Bay utilizing tray-held oysters from other Texas bays suggested that South Bay oysters had greater survival rates but those data were not statistically analyzed (Hofstetter et al. 1966, Hofstetter 1967). Previous studies designed to evaluate the re-establishment of oysters in Corpus Christi Bay have been inconclusive. One study suggested the possibility of successful re-establishment, while another resulted in total mortality of test oysters (Martinez 1963, 1964). However, oysters in these studies were not exposed to similar conditions prior to study initiation. If Eastern oysters from Corpus Christi Bay are unable to survive in harvestable numbers because of environmental conditions present in the bay, it may be possible to repopulate the bay by stocking animals preadapted to conditions similar to those found in Corpus Christi Bay. Oysters in the lower Laguna Madre and South Bay tolerate salinities > 40 o/oo (Breuer 1962). These oysters may be able to adapt to conditions in Corpus Christi Bay.

The purpose of this study was to determine if Eastern oysters from other Texas bay systems, especially South Bay, might be suitable for restocking Corpus Christi Bay. Specific study objectives were to compare growth and survival of Eastern oysters from six Texas bay systems held in Corpus Christi Bay.

## MATERIALS AND METHODS

Eastern oysters were collected by dredge or hand from Galveston, Matagorda, San Antonio, Aransas, Nueces (Corpus Christi Bay system) and South Bays during summer 1988. About 200 specimens were collected from each bay system and transported on ice to the Perry R. Bass Marine Fisheries Research Station near Palacios, Texas. Oysters were allowed to warm to room temperature before being placed in a 4,000-l saltwater tank system where they remained up to 3 weeks. Unfiltered Matagorda Bay water, flowed through the system using a submersible pump (Model A-01, Fritz Aquaculture, Dallas, Texas), provided a food source. Temperature and salinity were monitored daily using a mercury glass thermometer and a salinity-conductivity-temperature meter (Model 33, Yellow Springs Instrument Company, Yellow Springs, Ohio), respectively.

After acclimatization, about 50 Eastern oysters from each bay were transported on ice to Corpus Christi Bay on 29 June 1988. The remaining oysters from each bay were retained in the holding system. Each individual oyster was placed in a numbered 10 x 15-mm nylon bag of 6-mm mesh. Five oysters from each bay system were then placed into each of five replicate cages. The cages consisted of 35 X 35-mm plastic milk crates with a 6-mm mesh cover. The study site was in close proximity to historical Alto Vista reef (Figure 1). The cages were secured to the 4600 building pier on Ocean Drive, Corpus Christi, Texas. Each cage was suspended about 1 m from the substrate to reduce sediment buildup and allow for tidal fluctuations. Following a 2-week acclimation period, all dead oysters were replaced from the surplus in the holding system. The study was initiated 12 July 1988.

Growth and mortality were monitored every 12-21 days for 153 days. Total shell length for each individual was determined by measuring from the beak of the right valve to the furthest point on the shell margin. Temperature and salinity were measured every sampling period.

A repeated measures analysis of variance (ANOVA) was used to test for differences in total shell length of oysters among bay systems (Sokal and Rohlf 1981). During the course of the study, one cage was lost at day 56 due to vandalism and two were lost at day 72 from the effects of hurricane Gilbert on 14-16 September 1988. Therefore, two data sets were analyzed to maximize statistical power. One data set included all cages to day 56 (first cage lost), while the second set analyzed data from the two cages that remained at study termination.

Total shell length was regressed against time and the heterogeneity of slopes was tested to determine if differences existed in shell growth rates among bay systems (Sokal and Rohlf 1981). Percent survival was calculated for each cage and bay. Log likelihood ratio G test using maximum likelihood ratio statistics was used to test for significance in survival between bay systems (SAS Institute Incorporated 1985). Log likelihood ratio of all pairwise comparisons was performed. G values of all comparisons were compared to Table 15 of Sokal and Rohlf (1981); G values > 8.6 were significant. For all analyses, probability of 0.05 was used as test significance criteria.

At the end of the study all surviving oysters were examined for Perkinsus marinus using methods of Mackin (1962). A Spearman rank correlation (r) analysis was performed to determine if there was a relationship between the magnitude of Perkinsus marinus infection and growth or survival (Sokal and Rohlf 1981).

## RESULTS

No difference in mean total shell length of Eastern oysters from six Texas Bay systems were found in four cages from study initiation through day 56 ( $F_{5,56}=0.56$ ,  $P=0.73$ ) or in two cages from study initiation through termination ( $F_{5,17}=0.69$ ,  $P=0.64$ ) (Tables 1 and 2). Appendix A contains mean ( $\pm$  SD) shell length presented by cage and day with univariate F-statistics (Table A.1) and raw data (Table A.2).

Growth rates for South Bay were significantly different from San Antonio Bay oysters ( $t=2.05$ ,  $P=0.04$ ) at day 56 (day first cage was lost) (Table 1, Figure 2) and from Aransas Bay ( $t=0.64$ ,  $P=0.01$ ) at 153 days (Table 2, Figure 3). Growth comparisons (Y-intercepts) at study termination further suggested that Matagorda Bay oysters were larger than Galveston ( $t=-2.11$ ,  $P=0.04$ ) and South Bays ( $t=-1.97$ ,  $P=.04$ ) throughout the study (Table 2). However, growth rates were similar for Matagorda, South and Galveston Bay oysters (Table 2).

There were no statistically significant differences in survival among bay systems at day 56 ( $X^2 = 8.68$ ,  $df= 5$ ,  $P = 0.12$ ) or day 153 ( $X^2 = 7.23$ ,  $df=5$ ,  $P= 0.30$ ). However, at day 42, South Bay oysters exhibited significantly higher survival than Galveston, Nueces and San Antonio Bays ( $X^2 = 12.96$ ,  $df=5$ ,  $P= 0.02$ ) (Table 3). Survivals by cage are presented in Appendix B Table B.1.

The incidence of Perkinsus marinus infection in oyster mantle and rectum tissue examined at study termination was 100% in each bay system. No significant correlation was observed between the concentration in the mantle on growth ( $r=0.35$ ,  $P=0.49$ ) or survival ( $r=-0.11$ ,  $P=0.84$ ). Similarly, no significant correlations were found between concentrations in the rectum on growth ( $r= -0.20$ ,  $P=0.70$ ) or survival ( $r=-0.27$ ,  $P=0.60$ ).

Mean surface water temperatures and salinities were 30.7 C and 36.2 o/oo up to day 56, and 25.8 C, and 36.4 o/oo up to day 153. Surface water temperatures and salinities ranged 12.0-33.3 C and 30-40 o/oo, respectively (Table 4).

## DISCUSSION

Results of this study suggest there is no distinct advantage in stocking Eastern oysters from bay systems other than Corpus Christi (Nueces Bay). However, sample size was small and the study was conducted for only 6 months,

which may not be sufficient to distinguish all the environmental effects on the oysters (e.g. winter temperatures). Fouling organisms such as barnacles, sea squirts, and mussels may not have had time to attach to the oysters. These organisms could affect survival by competing for food and possibly cause a differential mortality rate among the different groups of oysters. Although all survivors were infected with Perkinsus marinus, oysters were not examined prior to study initiation. Therefore, any observed mortalities cannot be positively attributed to Perkinsus infection.

The differences in survival at day 42 could be attributed to handling procedures. Oysters from all bay systems except South Bay were collected by dredging. South Bay oysters were hand collected. Those oysters collected by dredging may have been under more stress thus causing a higher initial mortality rate.



## LITERATURE CITED

- Breuer, J. P. 1962. An ecological survey of the lower Laguna Madre. Publications of the Institute of Marine Science, University of Texas. 8:153-183.
- Hofstetter, R. P., T. L. Heffernan, and B. D. King III. 1966. Oyster mortality studies along the Texas coast. Coastal Fisheries Project Report 1965. Texas Parks and Wildlife Department. Austin, Texas.
- Hofstetter, R. P. 1967. Oyster (Crassostrea virginica) studies along the Texas coast. Coastal Fisheries Project Report 1966. Texas Parks and Wildlife Department. Austin, Texas.
- Longwell, A. C., and S. S. Stiles. 1970. The genetic system and breeding potential of the commercial American oyster. Endeavour. 29(107):94-99.
- Mackin, J. G. 1962. Report on studies of oyster parasites in the Aransas Bay and Galveston Bay areas of Texas. Coastal Fisheries Branch Project Report 1962. Texas Parks and Wildlife Department. Austin, Texas.
- Malinowski, S. and R.B. Whitlatch. 1988. A theoretical evaluation of shellfish resource management. Journal of Shellfish Research. 7(1):95-100.
- Martinez, R. 1963. Study of oyster population and experimental planting in Corpus Christi Bay. Annual Report of the Texas Parks and Wildlife Department 1963. Austin, Texas.
- Martinez, R. J. 1964. Analysis of populations of sports and commercial finfish and of factors which affect these populations in the coastal bays of Texas - Population studies of the sports and commercial fin-fish species of the Corpus Christi Bay system. Coastal Fisheries Project Report 1964. Texas Parks and Wildlife Department. Austin, Texas.
- Quast, W. D., M. A. Johns, D. E. Pitts, Jr., G. C. Matlock, and J. E. Clark. 1988. Texas Oyster Management Plan Source Document. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- SAS Institute Incorporated. 1985. SAS/STAT Guide for personal computers. Cary, North Carolina.
- Sokal, R. R., and F. J. Rohlf. 1981. Biometry. W. H. Freeman and Company. San Francisco, California.
- Yonge, C. M. 1960. Oysters. Collins Publishing. London.

Table 1. Mean ( $\pm$  SD) total shell lengths (mm), growth rates (slope), and percent survival of Eastern oysters from six Texas bays held in four cages in Corpus Christi Bay for 56 days. Slope and Y-intercept values followed by a different letter in a column are significantly different ( $P < 0.05$ ).

Bay	Mean total length		Slope	Y-intercept	% Survival N
	Day 0	Day 56			
Aransas	63.66 $\pm$ 13.37	70.40 $\pm$ 11.52	2.12 AB	59.71 A	75 15
Galveston	65.70 $\pm$ 14.31	71.55 $\pm$ 12.66	1.86 AB	62.45 A	45 9
Matagorda	67.69 $\pm$ 11.29	73.61 $\pm$ 10.03	1.99 AB	64.05 A	65 13
Nueces	67.64 $\pm$ 6.26	75.09 $\pm$ 7.12	2.43 AB	63.16 A	55 11
San Antonio	67.77 $\pm$ 11.06	72.92 $\pm$ 8.21	1.76 B	64.70 A	65 13
South	65.32 $\pm$ 12.43	78.12 $\pm$ 13.17	4.14 A	58.17 A	95 19

Table 2. Mean ( $\pm$  SD) total shell lengths (mm), growth rate (slope) and percent survival of Eastern oysters from six Texas bays held in two cages in Corpus Christi Bay for 153 days. Slope and Y-intercept values followed by a different letter in a column are significantly different ( $P < 0.05$ ).

Bay	Mean total length		Slope	Y-intercept	% Survival	N
	Day 0	Day 153				
Aransas	64.00 $\pm$ 15.40	70.67 $\pm$ 12.50	0.67 A	64.85 AB	30	3
Galveston	60.67 $\pm$ 18.50	70.67 $\pm$ 14.01	0.95 AB	59.07 B	30	3
Matagorda	70.20 $\pm$ 14.92	76.20 $\pm$ 13.77	0.67 AB	70.93 A	50	5
Nueces	64.60 $\pm$ 5.98	75.40 $\pm$ 10.92	1.11 AB	63.56 AB	50	5
San Antonio	68.00 $\pm$ 6.38	74.25 $\pm$ 4.19	0.73 AB	67.54 AB	40	4
South	63.33 $\pm$ 8.60	79.89 $\pm$ 10.34	1.78 B	62.48 B	90	9

Table 3. Survival in Eastern oysters maintained in Corpus Christi Bay, Texas by sampling day and chi-square approximations of log likelihood ratio tests between bay systems. A pair-wise comparison of the log likelihood (G) ratio test was performed on day 42, counts followed by unlike letters are significantly different ( $P < 0.01$ ).

Bay	0	Number of oysters surviving at sample day										153
		21	42 <sup>A</sup>	56 <sup>B</sup>	72 <sup>C</sup>	92	111	126	141			
Aransas	25	20	19 <sup>XY</sup>	14	6	4	4	3	3	3	3	3
Galveston	25	14	11 <sup>Y</sup>	9	3	3	3	3	3	3	3	3
Matagorda	25	16	16 <sup>XY</sup>	13	8	6	6	5	5	5	5	5
Nueces	25	17	14 <sup>Y</sup>	11	5	5	5	5	5	5	5	5
San Antonio	25	15	14 <sup>Y</sup>	13	7	7	6	4	4	4	4	4
South	25	24	24 <sup>X</sup>	18	10	10	10	9	9	9	9	9
chi-square		9.70	12.96	8.68	5.53	3.80	2.64	7.23	7.23	7.23	7.23	7.23

A Initial N = 25.

B Initial N = 20 due to 1 cage loss.

C Initial N = 10 due to 3 cage losses.

Table 4. Surface water temperatures and salinities of Corpus Christi Bay at the cage study location.

Day	Temperature (C)	Salinity (o/oo)
0	31.6	36
21	31.4	35
42	33.3	40
56	26.3	34
72	28.9	30
92	23.6	38
111	25.0	38
126	25.3	38
141	16.0	35
153	12.0	40

Figure 1. Study location in Corpus Christi Bay.






Figure 2. Mean total length (mm) of Eastern oysters held in Corpus Christi Bay for 56 days.





Figure 3. Mean total length (mm) of Eastern oysters held in Corpus Christi Bay for 153 days.



Appendix A: Total length data from Eastern oysters held in Corpus Christi Bay.



Table A.1. Mean ( $\pm$  SD) total shell lengths (mm) and sample size (N) by cage of Eastern oysters from six Texas bays held in Corpus Christi Bay including F-statistics for all bays and cages by sample day.

Day	Bay	Cage				
		1	2	3	4	5
0	Aransas (N)	67.50 $\pm$ 15.26 (5)	66.33 $\pm$ 12.74 (5)	65.50 $\pm$ 17.69 (5)	63.67 $\pm$ 5.51 (5)	62.75 $\pm$ 10.14 (5)
	Galveston (N)	51.50 $\pm$ 13.43 (5)	73.25 $\pm$ 7.32 (5)	71.75 $\pm$ 11.84 (5)	66.67 $\pm$ 10.12 (5)	46.00 $\pm$ 0.00 (5)
	Matagorda (N)	57.50 $\pm$ 16.26 (5)	74.75 $\pm$ 9.53 (5)	71.25 $\pm$ 21.75 (5)	80.00 $\pm$ 9.90 (5)	71.33 $\pm$ 6.66 (5)
	Nueces (N)	66.00 $\pm$ 3.46 (5)	65.25 $\pm$ 6.08 (5)	66.60 $\pm$ 7.02 (5)	55.50 $\pm$ 9.19 (5)	70.25 $\pm$ 6.40 (5)
	San Antonio (N)	75.20 $\pm$ 10.06 (5)	67.00 $\pm$ 10.44 (5)	62.40 $\pm$ 10.14 (5)	79.25 $\pm$ 5.85 (5)	69.25 $\pm$ 9.91 (5)
	South (N)	56.75 $\pm$ 9.64 (5)	62.00 $\pm$ 9.13 (5)	65.00 $\pm$ 9.62 (5)	68.80 $\pm$ 7.05 (5)	75.50 $\pm$ 17.44 (5)
	F-statistic	$F_{5,56}=0.27, P = 0.9569$				
21	Aransas (N)	66.40 $\pm$ 13.85 (5)	71.50 $\pm$ 7.59 (5)	64.80 $\pm$ 16.48 (5)	65.50 $\pm$ 6.45 (5)	66.50 $\pm$ 4.95 (5)
	Galveston (N)	56.00 $\pm$ 11.31 (5)	71.67 $\pm$ 8.08 (5)	74.25 $\pm$ 13.05 (5)	67.00 $\pm$ 8.54 (5)	66.00 $\pm$ 19.80 (5)
	Matagorda (N)	66.75 $\pm$ 11.35 (5)	76.75 $\pm$ 8.81 (5)	65.67 $\pm$ 12.50 (5)	75.67 $\pm$ 11.01 (5)	71.50 $\pm$ 6.36 (5)
	Nueces (N)	70.50 $\pm$ 13.08 (5)	67.50 $\pm$ 5.97 (5)	71.33 $\pm$ 6.66 (5)	70.00 $\pm$ 17.78 (5)	74.33 $\pm$ 5.03 (5)

Table A.1. (Cont'd.)

Day	Bay	Cage				
		1	2	3	4	5
	San Antonio (N)	76.00 ± 10.07 (5)	62.50 ± 3.54 (5)	62.33 ± 7.51 (5)	86.00 ± 2.83 (5)	74.00 ± 7.00 (5)
	South (N)	66.20 ± 11.71 (5)	65.20 ± 6.57 (5)	72.40 ± 10.53 (5)	73.60 ± 9.21 (5)	88.00 ± 14.26 (5)
	F-statistic	F <sub>5,56</sub> =0.51, P = 0.7696				
42	Aransas (N)	64.60 ± 3.46 (5)	71.00 ± 8.89 (4)	67.40 ± 6.39 (5)	67.25 ± 6.60 (4)	71.50 ± 4.95 (2)
	Galveston (N)	57.50 ± 3.43 (2)	77.00 ± 0.00 (3)	75.50 ± 1.27 (4)	73.50 ± 7.78 (3)	67.00 ± 8.38 (2)
	Matagorda (N)	68.25 ± 1.79 (4)	78.50 ± 9.95 (4)	67.00 ± 2.12 (3)	76.33 ± 11.01 (3)	77.00 ± 7.07 (2)
	Nueces (N)	69.00 ± 12.29 (4)	72.67 ± 4.04 (4)	72.50 ± 9.19 (3)	72.33 ± 17.62 (3)	77.33 ± 4.04 (3)
	San Antonio (N)	77.60 ± 9.39 (5)	65.00 ± 0.00 (2)	65.33 ± 4.16 (3)	88.00 ± 0.00 (1)	76.00 ± 3.46 (3)
	South (N)	69.40 ± 12.22 (5)	67.60 ± 8.08 (5)	77.20 ± 9.68 (5)	77.20 ± 10.52 (5)	88.75 ± 14.68 (4)
	F-statistic	F <sub>5,56</sub> =0.82, P = 0.5419				

Table A.1. (Cont'd.)

Day	Bay	Cage				
		1	2	3	4	5
56	Aransas (N)	70.50 ± 8.48 (5)	72.33 ± 10.08 (3)	71.60 ± 7.51 (5)	ND	74.00 ± 15.39 (2)
	Galveston (N)	57.50 ± 13.43 (2)	77.00 ± 0.00 (1)	77.50 ± 8.10 (4)	ND	71.00 ± 18.38 (2)
	Matagorda (N)	70.00 ± 10.42 (4)	77.75 ± 9.29 (4)	68.00 ± 12.17 (3)	ND	81.00 ± 1.41 (2)
	Nueces (N)	69.67 ± 11.59 (3)	76.33 ± 2.08 (3)	78.00 ± 8.48 (2)	ND	77.33 ± 4.40 (3)
	San Antonio (N)	78.00 ± 9.35 (5)	66.50 ± 2.12 (2)	65.67 ± 4.04 (3)	ND	76.00 ± 3.46 (3)
South (N)	72.00 ± 12.41 (5)	69.60 ± 7.30 (5)	81.60 ± 11.82 (5)	ND	87.33 ± 4.16 (4)	
	F-statistic	F <sub>5,56</sub> = 1.36, P = 0.2517				
72	Aransas (N)	70.67 ± 9.71 (4)	73.33 ± 6.81 (3)	ND	ND	ND
	Galveston (N)	57.50 ± 13.43 (2)	77.00 ± 0.00 (1)	ND	ND	ND
	Matagorda (N)	70.00 ± 10.42 (4)	77.75 ± 9.29 (4)	ND	ND	ND

Table A.1. (Cont'd.)

Day	Bay	Cage				
		1	2	3	4	5
	Nueces (N)	63.00 ± 1.41 (3)	76.67 ± 2.52 (3)	ND	ND	ND
	San Antonio (N)	79.20 ± 7.26 (5)	67.50 ± 0.71 (2)	ND	ND	ND
	South (N)	73.80 ± 13.29 (5)	71.40 ± 7.44 (5)	ND	ND	ND
	F-statistic	F <sub>5,17</sub> = 0.79, P = 0.5734				
92	Aransas (N)	70.50 ± 14.85 (3)	76.00 ± 8.49 (3)	ND	ND	ND
	Galveston (N)	58.50 ± 14.85 (2)	74.00 ± 0.00 (1)	ND	ND	ND
	Matagorda (N)	65.67 ± 9.29 (4)	85.00 ± 9.16 (4)	ND	ND	ND
	Nueces (N)	63.00 ± 1.41 (2)	76.67 ± 2.52 (3)	ND	ND	ND
	San Antonio (N)	79.20 ± 7.26 (5)	67.50 ± 0.71 (2)	ND	ND	ND
	South (N)	73.80 ± 13.29 (5)	71.40 ± 7.44 (5)	ND	ND	ND
	F-statistic	F <sub>5,17</sub> = 0.93, P = 0.4848				

Table A.1. (Cont'd.)

Day	Bay	Cage				
		1	2	3	4	5
111	Aransas (N)	71.50 ± 14.85 (2)	76.00 ± 8.49 (2)	ND	ND	ND
	Galveston (N)	58.50 ± 14.85 (2)	76.00 ± 0.00 (1)	ND	ND	ND
	Matagorda (N)	67.33 ± 8.14 (3)	86.00 ± 9.16 (3)	ND	ND	ND
	Nueces (N)	63.00 ± 1.41 (2)	79.33 ± 4.04 (3)	ND	ND	ND
	San Antonio (N)	80.60 ± 6.15 (5)	68.00 ± 0.00 (2)	ND	ND	ND
	South (N)	78.00 ± 17.10 (5)	71.60 ± 9.13 (5)	ND	ND	ND
F-statistic		F <sub>5,17</sub> = 0.94, P = 0.4815				
126	Aransas (N)	62.00 ± 0.00 (1)	75.50 ± 7.78 (2)	ND	ND	ND
	Galveston (N)	60.00 ± 1.14 (2)	77.00 ± 0.00 (1)	ND	ND	ND
	Matagorda (N)	66.00 ± 11.31 (3)	87.00 ± 8.18 (3)	ND	ND	ND
	Nueces (N)	63.50 ± 2.12 (2)	78.33 ± 3.51 (3)	ND	ND	ND

Table A.1. (Cont'd.)

Day	Bay	Cage				
		1	2	3	4	5
	San Antonio (N)	77.67 ± 2.08 (5)	69.00 ± 0.00 (1)	ND	ND	ND
	South (N)	85.50 ± 11.96 (5)	72.60 ± 8.20 (5)	ND	ND	ND
	F-statistic	F <sub>5,17</sub> = 1.22, P = 0.3406				
141	Aransas (N)	59.00 ± 0.00 (1)	76.50 ± 7.78 (2)	ND	ND	ND
	Galveston (N)	66.00 ± 11.31 (2)	84.00 ± 0.00 (1)	ND	ND	ND
	Matagorda (N)	66.50 ± 16.26 (3)	83.00 ± 5.00 (3)	ND	ND	ND
	Nueces (N)	63.50 ± 6.36 (2)	81.67 ± 8.33 (3)	ND	ND	ND
	San Antonio (N)	74.33 ± 5.13 (3)	68.00 ± 0.00 (1)	ND	ND	ND
	South (N)	87.00 ± 12.91 (4)	75.20 ± 6.76 (5)	ND	ND	ND
	F-statistic	F <sub>5,17</sub> = 1.29, P = 0.3126				



Table A.1. (Cont'd.)

Day	Bay	Cage				
		1	2	3	4	5
153	Aransas (N)	58.00 ± 0.00 (1)	77.00 ± 8.48 (2)	ND	ND	ND
	Galveston (N)	65.00 ± 14.14 (2)	82.00 ± 0.00 (1)	ND	ND	ND
	Matagorda (N)	64.50 ± 13.43 (2)	84.00 ± 7.81 (3)	ND	ND	ND
	Nueces (N)	64.50 ± 3.54 (2)	82.67 ± 5.86 (3)	ND	ND	ND
	San Antonio (N)	75.67 ± 3.79 (3)	70.00 ± 0.00 (1)	ND	ND	ND
	South (N)	90.00 ± 6.24 (4)	73.80 ± 8.44 (5)	ND	ND	ND
	F-statistic	F <sub>5,17</sub> = 1.40, P = 0.2729				

ND No data available, cage lost.



Table A.2. (Cont'd)

Bay	Cage	Oyster bag number	Length at sample day														
			0	21	42	56	72	92	111	126	141	153					
Galveston	2	1537	78	77													
	2	1538	86	85													
	2	1539	63	63	64												
	2	1540	73	73	72												
	2	1541	79	79	77	77	77	74	76	77	84	82					
	3	1567	82	88													
	3	1568	78	79	79	79											
	3	1569	84	88	88	88											
	3	1570	68	73	74	74											
	3	1571	57	57	61	69											
	4	2926	73	75	79												
	4	2927	55	58	58												
	4	2928	62	68	68												
	4	2929	72	75													
	4	2930	84	84													
	5	2956	73	75													
	5	2957	55	56													
5	2958	46	52	54	58												
5	2959	77	80	80	84												
5	2960	54	55														
Maragorda	1	1512	98	98													
	1	1513	46	51	52	56	56	55	58	58	55	55					
	1	1514	69	70	72	73	73	72	73	74	78	74					
	1	1515	68	68	69	70	70	70	71	71	71						
	1	1516	75	78	80	81	81	81	81	81	81						
	2	1542	83	83	86	86	86	93	94	94	88	88					
	2	1543	82	84	86	83	83	87	88	88	83	89					
	2	1544	72	74													
	2	1545	71	75	77	77	77	75	76	78	78	75					
	2	1546	63	65	65	65	65	65	65	65	65	65					





Table A.2. (Cont'd)

Bay	Cage	Oyster bag number	Length at sample day															
			0	21	42	56	72	92	111	126	141	153						
South	5	2983	90	95														
	5	2984	76	84	84	84												
San Antonio	5	2985	63	92	92	92												
	1	1517	73	74	76	78	78	79	80	80	80	80	80	80	80	80	80	80
	1	1518	74	74	77	77	77	74	76	76	76	76	76	76	73	73	73	73
	1	1519	63	63	64	64	70	75	76	76	77	77	77	77	70	70	70	74
	1	1520	75	78	81	81	81	79	80	80	80	80	80	80				
	1	1521	91	91	90	90	90	91	91	91	91	91	91	91				
	2	1547	79	79														
	2	1548	65	65														
	2	1549	60	60	65	68	68	63	65	65	68	68	68	68	68	68	68	68
	2	1550	62	65	65	65	67	67	67	68	69	69	69	69	68	68	68	68
2	1551	86	89															
3	1578	64	70	70	70													
3	1579	78	78															
3	1580	50	55	62	62													
3	1581	60	62	64	65													
3	1582	60	61															
4	2936	81	84	84														
4	2937	78	78															
4	2938	86	87															
4	2939	72	72															
4	2940	86	88	88														
5	2966	56	66	72	72													
5	2967	79	79	78	78													
5	2968	68	68															
5	2969	74	77	78	78													
5	2970	88	88															



Appendix B. Survival data from Eastern oysters held in Corpus Christi Bay.

Table B.1. Survival (number) by cage of Eastern Oysters held in Corpus Christi Bay.

Bay	Cage	Number of live oysters at sample day										
		0	21	42	56	72	92	111	126	141	153	
Aransas	1	5	5	5	5	4	3	2	1	1	1	1
	2	5	5	4	3	3	3	2	2	2	2	2
	3	5	5	5	5	ND	ND	ND	ND	ND	ND	ND
	4	5	5	4	ND	ND	ND	ND	ND	ND	ND	ND
	5	5	5	2	2	ND	ND	ND	ND	ND	ND	ND
Galveston	1	5	5	2	2	2	2	2	2	2	2	2
	2	5	5	3	1	1	1	1	1	1	1	1
	3	5	5	4	4	ND	ND	ND	ND	ND	ND	ND
	4	5	5	3	ND	ND	ND	ND	ND	ND	ND	ND
	5	5	5	2	2	ND	ND	ND	ND	ND	ND	ND
Matagorda	1	5	5	4	4	4	4	3	3	3	3	2
	2	5	5	4	4	4	4	3	3	3	3	3
	3	5	5	3	3	ND	ND	ND	ND	ND	ND	ND



Table B.1. (Cont'd.)

Bay	Cage	Number of live oysters at sample day										
		0	21	42	56	72	92	111	126	141	153	
South	4	5	5	5	ND	ND	ND	ND	ND	ND	ND	ND
	5	5	5	4	4	ND	ND	ND	ND	ND	ND	ND
San Antonio	1	5	5	5	5	5	5	5	5	3	3	3
	2	5	5	2	2	2	2	2	1	1	1	1
	3	5	5	3	3	ND	ND	ND	ND	ND	ND	ND
	4	5	5	2	ND	ND	ND	ND	ND	ND	ND	ND
	5	5	5	3	3	ND	ND	ND	ND	ND	ND	ND

ND No data available, cage lost.

**PWD-RP-3400-341-1/91**

Dispersal of this publication conforms with Texas State Documents Depository Law, and it is available at Texas State Publications Clearinghouse and Texas Depository Libraries.