Trends in Abundance and Size of Adult White Shrimp in Texas Gulf Waters
≤18.3 m During
May-July 1970-82

by
Billy E. Fuls
Lawrence W. McEachron
and
C.E. Bryan

Management Data Series No. 42 1990



TEXAS
PARKS & WILDLIFE
DEPARTMENT
FISHERIES DIVISION

4200 Smith School Road Austin, Texas 78744

TRENDS IN ABUNDANCE AND SIZE OF ADULT WHITE SHRIMP IN TEXAS GULF WATERS ≤18.3 m DURING MAY-JULY 1970-82

by

Billy E. Fuls, Lawrence W. McEachron and C. E. Bryan

MANAGEMENT DATA SERIES No. 42 1990

Texas Parks and Wildlife Department
Fisheries Division
Coastal Fisheries Branch
4200 Smith School Road
Austin, Texas 78744

ACKNOWLEDGEMENTS

We would like to thank the various crews aboard the R/V WESTERN GULF for the collection of shrimp samples from 1970 to 1982. Appreciation is extended to Gary Matlock, Jerry Clark, Gary Saul, Tom Heffernan, Ed Hegen, Lynn Benefield, and Terry Cody for reviewing the manuscript. Thanks is extended to Al Green and Jeff Doerzbacher for their statistical comments and review. This study was conducted with partial funding from the U. S. Department of Commerce; National Oceanic and Atmospheric Administration, National Marine Fisheries Service, under P.L. 88-309.

ABSTRACT

Analysis of catch rates (no./h) and mean length of white shrimp ($\underline{Penaeus}$ setiferus) collected in gulf waters (≤ 18.3 m deep) off the central Texas coast during May-July 1970-82 revealed a significant difference among years in mean length but not in catch rate. Linear regression analysis revealed a negative linear trend in mean length among years, depicting a decrease in size of white shrimp through time.

INTRODUCTION

Penaeid shrimp constitute Texas' most valuable commercial fishery. White shrimp (<u>Penaeus setiferus</u>) comprised 25% (by weight) of the annual mean (37.4 million Kg) combined reported commercial shrimp landings from Texas bays and the gulf during 1977-86 (Osburn et al. 1987). White shrimp comprised 53% of the reported landings from bays and 22% from the gulf. During 1986, reported commercial gulf landings of white shrimp were greatest (62%) during August-November; 21% were landed during May-July.

In 1981, implementation of the Gulf of Mexico Fishery Management Council's (GMFMC) shrimp management plan (Van Lopik et al. 1980) closed the Exclusive Economic Zone (EEZ) off Texas to shrimping in conjunction with the annual closure (typically 1 June-15 July) of the Texas Territorial Sea (TTS). However, it remained legal to catch white shrimp during daylight hours in the TTS during the Texas closure in water ≤ 7.3 -m deep.

The major white shrimp fishing grounds within the Texas gulf are at depths ≤18.3 m (Hildebrand 1954, Griffin et al. 1974). Overall mean length of white shrimp within the TTS is greatest from May-August then declines from September through December as emigrating young shrimp enter the gulf from Texas bays (Bryan et al. 1978). Mature white shrimp are found within shallow gulf waters mainly from April through August with peak spawning during June (Lindner and Anderson 1956, Perez-Farfante 1969, Bryan and Cody 1975, Chamberlain and Lawrence 1983), which parallels the annual closure of Texas gulf waters.

There has been concern whether historical to present-day white shrimp stock abundance and average shrimp size has remained relatively stable through time. Caillouet et al. (1980) observed that reported commercial white shrimp catches in Texas revealed a trend of decreasing size from 1959 to 1976. Rothschild and Brunenmeister (1984) and Brunenmeister (1984) observed that shrimp stocks, as indicated by CPUE (abundance and weight) of commercial shrimpers in the Gulf, were decreasing through time. The Texas Parks and Wildlife Department (TPWD) has monitored relative abundance and size of white shrimp since 1970 in gulf waters ≤ 18.3 m deep. The objective of the present study was to determine if there were differences and trends through time in relative abundance (no./h) and size (TL) of adult white shrimp off the central Texas gulf coast during May-July, 1970-82.

MATERIALS AND METHODS

White shrimp were collected with trawls aboard the R/V WESTERN GULF (a double-rigged, 21.9-m steel hull trawler) in gulf waters (\leq 18.3 m deep) off the central Texas coast (Figure 1) during May-July 1970-82. Samples were collected with trawls ranging in width from 12.2 to 14.3 m with stretched mesh ranging from 4.4 to 5.7 cm (Appendix A). Trawls were equipped with a tickler chain and spread by 0.9 x 2.1-m wooden doors. Day and night trawl samples were pooled because no significant differences have been found between day and night catches (Cody and Fuls 1986). Tow durations ranged from 5-60 minutes. All white shrimp were weighed en masse. Up to 50 randomly selected individuals were measured (TL to nearest mm).

Total number of shrimp in catches with >50 individuals was estimated using the following formula:

N = (T/S)(n)

where: N = total number of white shrimp in sample,

T = total weight of white shrimp in sample,

S = weight of white shrimp measured,n = number of white shrimp measured.

Catch rate (weighted to no./h) and total length data for white shrimp, associated with TPWD gulf trawl samples, were compiled by year (May-July) for analysis (Appendix A). Individual lengths were weighted for analysis by no./h for samples with all shrimp measured, and by number in sample not measured and no./h for samples with >50 shrimp. Data for 1975 were excluded from the analysis because of the small sampling effort and because samples were collected only in May.

SAS software programs (SAS Institute Inc. 1985) were used for all analyses. Using the SAS UNIVARIATE procedure, catch rates and lengths of white shrimp were found not to be normally distributed. Therefore, data were analyzed non-parametrically by ranking the data and testing for significant differences ($P \le 0.05$) among years using the SAS procedure GLM, for a one-way analysis of variance (AOV) with unequal sample sizes (Sokal and Rohlf 1981). The GT2-method for a one-way AOV with unequal sample size was used to determine which years were different (Sokal and Rohlf 1981).

Linear regression analysis involving multiple y at each x was performed regressing length on year to determine if a significant ($P \le 0.05$) linear trend existed (Sokal and Rohlf 1981).

Because only 1% of all white shrimp caught were <140 mm in total length it was assumed that white shrimp collected were maturing parental stock (Burkenroad 1939, Lindner and Anderson 1956). It was also assumed TPWD catches were proportional to shrimp populations throughout the sampling period and area despite various trawl sizes and tow durations.

RESULTS

The mean size of adult white shrimp off the central Texas coast decreased during 1970-82, but abundance remained relatively stable (Figures 2 and 3). Linear regression analysis revealed length was a significant function of year, however, the regression explained only 3% of the variation in mean length (Table 1). The regression further indicated a decrease in size through time (r = -0.18); overall size of adult white shrimp decreased about 5 mm from the period 1970-78 to 1979-82 (Figure 2). There was a significant difference in mean length (Figure 2, Table 1) but not in mean catch rate (Figure 3, Table 2) of white shrimp among years. Further analysis revealed mean length of white shrimp for 1971 significantly differed from all other years, whereas mean

length within grouped years of 70/72/74/76/77/78, 73/80/82, and 79/80/81 were similar among years within each group but differed among groups (Table 3).

DISCUSSION

Abundance of parental white shrimp in the gulf is dependent upon escapement from the bays. Results of this study indicated the magnitude of escapement (i.e. number of shrimp) remained constant during 1970-82. This was in spite of the fact number of white shrimp commercially caught in bays and number of bay and gulf licensed shrimp vessels increased during the same period (Cody et al. 1989). Reasons for this disparity would include: 1) recruitment to the parent stock came from other areas, or 2) fishing and natural mortality combined must have gone down, or 3) recruitment to the bays increased. The apparent reason is an increase in recruitment to bays since TPWD bag seine samples during 1978-82 indicated an overall increase in juvenile white shrimp catch rates (Cody et al. 1989). This finding conflicts with that of Rothschild and Brunenmeister (1984) who reported a decrease in Gulf-wide white shrimp stocks (in relation to abundance) through time.

If white shrimp mean length has decreased, as indicated by this study of parental stocks in the gulf and by TPWD bag seine samples of juveniles in bays (Cody et al. 1989), then the catch by weight off the central coast of Texas would show a decline if overall numbers of shrimp remained constant. The decrease of white shrimp length through time off the central Texas coast and in bays is supported by observations of Caillouet et al. (1980), and the decrease in weight is supported by Brunenmeister (1984) who reported a decrease in gulf-wide CPUE by weight.

It is documented that decrease in length of organisms in a fishery through time can be a function of increased fishing pressure (Cushing 1968, Gulland 1978, Ricker 1945, Royce 1972). However, it is not entirely clear why the length of white shrimp has decreased or why the numbers of recruits to the bays has increased. Perhaps the parental white shrimp stock has compensated for the increased fishing pressure on the overall shrimp population. If length of shrimp continues to decrease the minimum effect will be a decrease in the ex-vessel value of the catch because larger shrimp are more valuable. A more critical effect, however, would be an eventual collapse of the fishery since smaller shrimp produce fewer eggs (Chamberlain and Lawrence 1983).

Information presented in the present report on relative abundance and size of white shrimp should assist managers in constructing population models to manage the fishery. Additional analyses on white shrimp stocks coastwide within the Texas gulf during periods of major gulf recruitment and throughout the year should also give greater insight into abundance and size composition of white shrimp stocks within Texas waters. In 1985, the TPWD instituted a coastwide monthly gulf shrimp monitoring program, which will provide needed information to formulate more precise population models. To refine such models, and analyze long term effects on relative abundance and size of shrimp it is important that standardized gulf monitoring programs be maintained.

LITERATURE CITED

- Brunenmeister, S. R. 1984. Standardization of fishing effort and production models for brown, white and pink shrimp stocks fished in U.S. waters of the Gulf of Mexico. Pages 187-211. <u>In</u>: J. A. Gulland and B. J. Rothschild, editors. Penaeid shrimps-their biology and management. Fishing News Books Unlimited. England.
- Bryan, C. E., and T. J. Cody. 1975. White shrimp <u>Penaeus setiferus</u>
 (Linnaeus), spawning in the Gulf of Mexico off Texas 1973-75. Project 2202-4 Completion Report. Texas Parks and Wildlife Department, Coastal
 Fisheries Branch. Austin, Texas.
- Bryan, C. E., T. J. Cody, and K. W. Rice. 1978. Monitoring of shrimp populations and evaluation of the Texas shrimp laws in the Gulf of Mexico. Project 2-276-R, Part I, Completion Report. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- Burkenroad, M. D. 1939. Further observations on Penaeidae of the northern Gulf of Mexico. Bulletin Bingham Oceanography Collection 6.
- Caillouet, C. W., F. J. Patella, and W. b. Jackson. 1980. Trends toward decreasing size of brown shrimp, <u>Penaeus aztecus</u>, and white shrimp, <u>Penaeus setiferus</u>, in reported annual catches from Texas and Louisiana. Fisheries Bulletin. 77(4):985-990.
- Chamberlain, G. W., and A. L. Lawrence. 1983. Seasonal and spatial variations in reproductive activity and biochemical composition of <u>Penaeus setiferus</u> and <u>Penaeus aztecus</u> in the Gulf of Mexico. Texas A&M Sea Grant Publication (TAMU-SG-84-203). College Station, Texas.
- Cody, T. J., R. P. Campbell, P. C. Hammerschmidt, G. C. Matlock, C. E. Bryan, J. E. Clark, and L. S. Procarione. 1989. Texas shrimp fishery management plan; source document. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- Cushing, D. H. 1968. Fisheries biology, a study in population dynamics. University of Wisconsin Press. Madison, Wisconsin.
- Griffin, W. L., M. L. Cross, and G. W. Ryan. 1974. Seasonal and movement patterns in the Gulf of Mexico shrimp fishery. Department Technical Reports Number 74-4, Texas Agriculture Experiment Station, Texas A&M University. College Station, Texas.
- Gulland, J. A. 1978. Fish population dynamics. John Wiley and Sons. New York.

- Hildebrand, H. H. 1954. A study of the fauna of the brown shrimp (<u>Penaeus aztecus</u> Ives) grounds in the western Gulf of Mexico. Publications of the Institute of Marine Science. 3(2):233-366.
- Lindner, M. J., and W. W. Anderson. 1956. Growth, migrations, spawning and size distribution of shrimp <u>Penaeus setiferus</u>. Fisheries Bulletin. 56:555-645.
- Osburn, H. R., W. D. Quast, and C. L. Hamilton. 1987. Trends in Texas commercial fishery landings, 1977-1986. Management Data Series number 131. Texas Parks and Wildlife Department, Coastal Fisheries Branch. Austin, Texas.
- Perez-Farfante, I. 1969. Western Atlantic shrimps of the genus <u>Penaeus</u>. Fishery Bulletin. 67(3):461-591.
- Ricker, W. E. 1945. A method of estimating minimum size limits for obtaining maximum yield. Copeia. (2):85-94.
- Rothschild, B. J., and S. L. Brunenmeister. 1984. The dynamics and management of shrimp in the Northern Gulf of Mexico. Pages 145-172. <u>In</u>: J. A. Gulland and B. J. Rothschild, editors. Penaeid shrimps their biology and management. Fishing News Books Unlimited. England.
- Royce, W. F. 1972. Introduction to the Fishery sciences. Academic Press. New York.
- SAS Institute Incorporated. 1985. Procedures for personal computers, Version 6. Cary, North Carolina.
- Sokal, R. R., and F. J. Rohlf. 1981. Biometry, 2nd. Edition. W. H. Freeman and Company. San Francisco, California.
- Van Lopik, J. R., K. H. Drummond, and R. E. Condrey. 1980.

 Environmental impact statement and fishery management plan for the shrimp fishery of the Gulf of Mexico, United States waters. A regional management plan prepared for Gulf of Mexico Fishery Management Council, Tampa, Florida. Center for Wetland Resources, Louisiana State University. Baton Rouge, Louisiana.

Table 1. Summary of AOV table with regression for more than one value of Y [ranked TL (mm)] for each value of X (year) for white shrimp in TPWD gulf trawl collections within the 18.3-m depth zone during May-July 1970-82.

Source of variation	df	Sum of squares	F
Total	11452	12507365902	
Among years	11	7988589742	70.96
Linear regression	1	4125572956	106.80***
Deviations from regression	10	386301786	3.77***
Error	11441	117085067160	

 $[&]quot;P \le 0.001$

Table 2. Summary of AOV of ranked catch rates (no./h) for white shrimp in TPWD gulf trawl collections within the 18.3-m depth zone during May-July 1970-82. NS = not significant.

df	Sum of squares	F
196	629010	
11	53988	1.58 NS
185	575022	
	196 11	df squares 196 629010 11 53988

Table 3. Ranked mean TL (mm) by year for white shrimp in TPWD gulf trawl collections within the 18.3-m depth zone during May-July 1970-82. Means followed by the same letter in a column are similar (P >0.05).

Year	Unranked mean TL	Ranked mean TL
1971	174.5	7,731.7 A
1974	171.9	6,887.0 B
1972	171.0	6,791.8 B
1978	171.9	6,783.8 B
1977	170.9	6,507.1 B
1976	170.6	6,394.8 B
1970	170.2	6,278.6 B
1973	165.8	5,397.9 C
1982	167.1	5,314.8 C
1980	166.2	5,142.2 CD
1981	165.1	4,957.5 CD
1979	162.6	4,699.2 D

Figure 1. TPWD gulf trawl sample areas off the central Texas gulf coast during May-July 1970-82.






Figure 2. Mean length (\pm 2SE) of white shrimp in TPWD gulf trawl collections within the 18.3-m depth zone during May-July 1970-82.



YEAR

LENGTH(mm)

Figure 3. Mean catch rate (\pm 2SE) of white shrimp in TPWD gulf trawl collections within the 18.3-m depth zone during May-July 1970-82.



Appendix A. TPWD trawl catches within the gulf during May-July 1970-82.

Table A.1. Date, location, day or night sample, trawl size, mesh size, depth, tow duration, no./h and mean total length for individual sample collections of white shrimp caught within 18.3 m off the central Texas gulf coast by the TPWD during May-July 1970-82. ND = No data.

Year Date	Location	Day (D) or night (N)	Trawl size (m)	Mesh size (mm)	Water depth (m)	Tow duration (min)	No./h	Mean length (mm)
1970								
25 May	Cedar Bayou	Q	14.0	77	4.9	15	0	,
•	Cedar Bayou	Ω	14.0	77	14.6	09	31	170
08 Jun	Aransas Pass	Ω	14.0	77	7.3	15	380	171
	Aransas Pass	Q	14.0	38	12.8	2	0	
08 Jul	Aransas Pass	N	13.7	51	14.6	30	9	183
	Aransas Pass	Z	13.7	51	7.3	15	768	170
1971								
25 May	Aransas Pass	Q	13.7	57	5.5	15	12	156
		Z	13.7	57	11.0	15	80	175
		Z	13.7	57	16.4	15	16	174
08 Jun		Q	13.7	57	7.9	15	0	
		Z	13.7	57	14.6	15	84	171
21	Aransas Pass	N	13.7	57	7.3	15	9/	174
		Z	13.7	57	17.4	15	12	168
06 Jul	Aransas Pass	Q	13.7	57	12.8	15	0	
60		Z	13.7	57	14.6	15	∞	175
.	Aransas Pass	Z	13.7	57	7.3	15	8	174
17	Cedar Bayou	Z	13.7	57	14.6	15	7	196
	Gedar Bayou	Z	13.7	57	7.3	15	120	178
29	Aransas Pass	Q	13.7	57	11.9	15	26	174

Table A.1. (Cont'd.)

			Trawl	Mesh	Water	Tow		Mean
Year Date	Location	Day (D) or	size (m)	size (mm)	depth	duration (min)	N.	length
Nacc.	Pocacton	iit giit iii	7111	/ mmil /	/==/	/1117111/	110./11	/ IIIIII /
1972								
03 May	Aransas Pass	Q	Э.	51	18.3	15	32	165
19	Aransas Pass	Z	ω.	51	16.4	30	2	163
		Z	3.	51	7.3	15	∞	178
01 Jun	Aransas Pass	Z	Э.	51	16.4	15	∞	186
	Aransas Pass	Z	щ.	51	7.3	2	0	
60	Aransas Pass	Z	13.7	51	14.6	10	120	170
	Aransas Pass	Z	ж. :	51	9.1	10	138	173
27 Jul	Aransas Pass	Z	ω.	51	14.6	15	0	
28	Aransas Pass	N	3.	51	9.1	10	09	171
1973								
09 May	Cedar Bayou	Q	13.7	51	14.6	15	∞	167
•	Cedar Bayou	Q	13.7	51	7.3	10	282	149
30	Aransas Pass	Q	13.7	51	7.3	15	16	166
	Aransas Pass	Q	13.7	51	14.6	15	80	176
31	Aransas Pass	Q	13.7	51	7.3	15	0	
	Aransas Pass	Q	13.7	51	14.6	15	0	
25 Jun	Aransas Pass	Q	13.7	51	7.3	15	136	169
	Aransas Pass	Q	13.7	51	14.6	15	264	174
17 Jul	Aransas Pass	Q	13.7	54	11.9	15	188	173
	Aransas Pass	Q	13.7	54	13.7	15	32	175
	Aransas Pass	Q	13.7	54	7.3	15	52	175
1974								
11 Jun	Aransas Pass Aransas Pass Aransas Pass	000	13.7 13.7 13.7	51 51 51	7.3 11.0 14.6	15 15 15	79 79 79	150 169 172

Table A.1. (Cont'd.)

		·	Trawl	Mesh	Water	Tow		Mean
Year Date	Location	Day (D) or night (N)	size (m)	size (mm)	depth (m)	duration (min)	No./h	length (mm)
1974 (Cont'd.)	d.)							
18 Jun	Aransas Pass	Q	13.7	51	7.3	15	212	171
		Ω	13.7	51	11.0	15	7	181
	Aransas Pass	Q	13.7	51	14.6	15	0	
	Aransas Pass	N	13.7	51	14.6	15	0	
		Z	13.7	51	7.3	15	77	169
21	Aransas Pass	Q	13.7	51	14.6	15	8	176
	Aransas Pass	Q	13.7	51	11.0	15	79	174
	Aransas Pass	Q	13.7	51	7.3	15	48	171
27	Cedar Bayou	Q	13.7	51	7.3	15	28	178
	Cedar Bayon	D	13.7	51	11.0	15	116	174
	Cedar Bayon	D	13.7	51	14.6	15	140	170
15 Jul	Aransas Pass	Ω	13.7	51	7.3	15	0	
	Aransas Pass	Q	13.7	51	11.0	15	0	
	Aransas Pass	Ω	13.7	51	14.6	15	0	
22	Cedar Bayou	Ω	13.7	51	14.6	15	0	
	Cedar Bayou	Ω	13.7	51	11.0	15	0	
	Cedar Bayou	Q	13.7	51	7.3	15	7	181
1975								
77 May	Aransas Pass	C		51	•	15	0	
		Ω	13.7	51	11.0	15	84	160
		Q	•	51	•	15	156	168
		Z	•	51	•	15	16	160
Jun	ND							
Jul	ND							

Table A.1. (Cont'd.)

Year Date	Location	Day (D) or night (N)	Trawl size (m)	Mesh size (mm)	Water depth (m)	Tow duration (min)	No./h	Mean length (mm)
1976								
10 May	Aransas Pass	Z	13.7	77	14.6	30	26	164
11	Aransas Pass	Q	13.7	7 77	7.3	30	0	
	Aransas Pass	Q	13.7	7 7	•	30	2	165
23	Corpus Christi Pa	Pass N	13.7	7 7	11.0	30	26	163
54	Aransas Pass	Z	13.7	77	16.4	30	12	179
	Aransas Pass	Z	13.7	77	٠	15	32	170
09 Jun	Aransas Pass	Z	13.7	77	12.8	15	88	167
14	Aransas Pass	Q	13.7	7 7	•	30	80	185
	Aransas Pass	Q	•	77	14.6	30	96	169
16	Corpus Christi Pa	Pass N	•	777	•	30	12	178
21	Corpus Christi Pa	Pass N	•	77	•	15	16	180
	Corpus Christi Pa	Pass N	•	77	11.0	15	26	176
22	Corpus Christi Pa	Pass D		77	•	15	24	173
07 Jul	Aransas Pass	Z		51	•	15	204	167
80	Corpus Christi Pass		13.7	51	12.8	15	24	178
20	Aransas Pass	Q		51	•	30	28	177
	Aransas Pass	Q		51	•	30	07	177
21	Aransas Pass	Z		51		30	0	
27	Aransas Pass	Q	13.7	51	•	15	0	
	Aransas Pass	Q	•	51		15	80	180
	Aransas Pass	Q	13.7	51	14.6	15	12	187
	Aransas Pass	Q	13.7	51		10	0	
1977								
O May	Aransas Pass	<u>-</u>	13.7	5.1	7 3	30	0	170
25		aΩ	13.7	51	14.6	15	124	173

Table A.1. (Cont'd.)

			Trawl	Mesh	Water	Tow		Mean	ı
Year	Tocation	Day (D) or	size (m)	size (mm)	depth	duration (min)	No./h	length (mm)	1
Date	Tocac Toll	/::P::2::9:::	\\.						1
1977 (Cont'd.)	'd.)								
76 May	Aransas Pass	Z	13.7	51		30	89	170	
	ţ;	Pass	13.7	51	_	15	16	158	
27		Z	13.7	51	11.0	15	120	165	
31		Z	13.7	51	-	30	87	170	
06 Jun		Q	13.7	51		15	12	170	
	Christi	Pass N	13.7	51		30	∞	171	
29	Aransas Pass	Q	13.7	51		30	51	174	
		Q	13.7	51		15	0		
30		N	13.7	51		15	20	178	
,		Z	13.7	51		15	77	173	
13 Jul		D	13.7	51		30	112	174	
·		Q	13.7	51		30	0		フ
14		Z	13.7	51		30	0		
		Z	13.7	51	•	30	14	167	
15	Christi	Pass N	13.7	51	•	15	0		
19			13.7	51	•	15	128	171	
0									
19/8									
30 May	Aransas Pass	Q		51		30	0		
		Ω	3.	51		15	∞	170	
		Q		51	•	30	10	167	
		Q	щ	51	•	30	2	146	
31		Z	3.	51	•	15	87	170	
	Aransas Pass	Z	ж	51	•	15	52	163	
13 Jun	Aransas Pass	Z		51	•	30	9	176	
14	Aransas Pass	Z	13.7	51	11.0	30	9/	172	
	Aransas Pass	Q	•	51	•	15	0		

Table A.1. (Cont'd.)

			Trawl	Mesh	Water	Tow		Mean
Year Date	Location	Day (D) or night (N)	size (m)	size (mm)	depth (m)	duration (min)	No,/h	length (mm)
1978 (Cont.d)	7							
13100 (00115)								
14 Jun	Aransas Pass	Q		51	14.6	15	0	
29	Aransas Pass	Z	13.7	51	11.0	1.5	156	171
28	Aransas Pass	Z	•	51	14.6	15	72	173
	Aransas Pass	Q		51	7.3	15	16	184
	Aransas Pass	Q		51	14.6	15	77	166
14 Jul	Aransas Pass	Z		51	14.6	15	12	188
	Aransas Pass	Q		51	7.3	15	0	
	Aransas Pass	Q	•	51	14.6	15	7	181
15	Aransas Pass	Z	•	51	11.0	15	99	181
19	Aransas Pass	Q		51	7.3	15	0	
		Q	•	51	14.6	15	0	
1979								
15 May	Aransas Pass	Q		51	7.3	30	0	
16	Aransas Pass	Ω		51	9.1	30	0	
	Aransas Pass	Ω		51	12.8	30	0	
	Aransas Pass	Z		51	11.0	30	0	
	Aransas Pass	Z	13.7	51	14.6	30	∞	162
17	Aransas Pass	Z		51	7.3	30	0	
23	Aransas Pass	Q		51	16.4	30	14	167
24	Aransas Pass	Z		51	11.0	30	42	159
	Aransas Pass	Z		51	16.4	30	54	166
		Ω		51	7.3	30	77	160
	Aransas Pass	D		51	11.0	30	54	163
22 Jun	Aransas Pass	Z	13.7	51	14.6	30	0	

Table A.1. (Cont'd.)

Year Date	Location	Day (D) or night (N)	Trawl size (m)	Mesh size	Water depth	Tow duration (min)	No. /h	Mean length (mm)	
1979 (Cont'd.)	d.)								
30 Jul		۵ ډ	13.7	51	7.3	30	97	154	
1980	Aransas Pass	٩	13./	75	12.8	30	70	184	
20 May	Aransas Pass	z	13.7	51	12.8	30	14	157	
05 Jun		Q	13.7	51	7.3	15	77	191	
	Aransas Pass	Q	13.7	51		15	0		
1		Z	13.7	51		30	222	164	
25		Ω;	13.7	51		15	777	168	
08 Jul	Aransas Pass Aransas Pass	z z	13.7	51	16.4	30 15	87	168	21
60		iΩ	13.7	51		15	7	183	I
		Q	13.7	51		15	4	158	
24	Aransas Pass	Z	13.7	51		15	0		
25	Aransas Pass	Z	13.7	51		15	16	174	
1981									
06 May	Aransas Pass	Ω		51		10	99	164	
	Aransas Pass	Ω	\sim	51	11.0	10	0		
	Aransas Pass	Q		51		10	87	155	
	Aransas Pass	Ω	3	51		10	12	146	
12	Aransas Pass	Ω	33	51		10	9	185	
	Aransas Pass	Ω	3	51		10	9	142	
	Aransas Pass	Q	13.7	51	11.0	10	9	146	
	Aransas Pass	Q		51	7.3	10	0		

'able A.1. (Cont'd.)

Year		ay (D)	Trawl	Mesh	Water depth	Tow	;	Mean
Date	Location	night (N)	(m)	(mm)	(E)	(min)	No./h	(mm)
1981 (Cont'd.)	d.)							
19 May	Aransas Pass	Q	14.3	51	7.3	10	810	162
•	Aransas Pass	Ω	14.3	51	11.0	10	348	163
	Aransas Pass	Q	14.3	51		10	78	161
	Aransas Pass	Q	14.3	51	7.3	10	0	
21	Aransas Pass	Z	14.3	51	•	10	72	162
	Aransas Pass	Q	14.3	51	•	10	102	163
	Aransas Pass	Q	14.3	51		10	348	166
	Aransas Pass	Q	14.3	51	•	10	138	158
	Aransas Pass	Q	14.3	51	7.3	10	192	167
26	Aransas Pass	Q	14.3	51	•	10	168	170
٠	Aransas Pass	Q	14.3	51	•	10	198	165
	Aransas Pass	Q	14.3	51	•	10	258	165
	Aransas Pass	Q	14.3	51	•	10	36	173
27	Aransas Pass	Z	14.3	51	•	10	240	162
05 Jun	Aransas Pass	Q	14.3	51	7.3	10	99	168
	Aransas Pass	D		51	•	10	48	164
	Aransas Pass	Q	14.3	51	•	10	18	169
	Aransas Pass	Q	•	51	•	10	72	168
16	Aransas Pass	Q	14.3	51	7.3	10	12	176
	Aransas Pass	Ω	•	51	11.0	10	12	184
17	Aransas Pass	D	•	51	7.3	10	42	176
	Aransas Pass	Q	12.2	51	11.0	10	168	175
26	Aransas Pass	Q	•	51	•	10	09	169
	Aransas Pass	Q	•	51	•	10	282	170
08 Jul	Aransas Pass	Z	٠	51	7.3	10	30	165
60	Aransas Pass	Q	•	51	٠	10	30	147
	Aransas Pass	Q	•	51	11.0	10	36	174
30	Aransas Pass	Z	13.7	51	14.6	30	9	181

Table A.1. (Cont'd.)

Year		Day (D) or	Trawl size	Mesh size	Water depth	Tow duration		Mean length	
Date	Location	night (N)	(m)	(mm)	(m)	(min)	No./h	(mm)	
1982									
18 May	San Jose Island	Q	12.2	51	9.1	15	28	166	
	San Jose Island	Ω	12.2	51	9.1	15	77	160	
	San Jose Island	Ω	12.2	51	17.4	15	36	165	
	San Jose Island	Ω	12.2	51	18.3	15	88	173	
	San Jose Island	Ω	12.2	51	17.9	15	152	165	
19	Aransas Pass	Ω	12.2	51	17.9	15	72	167	
	Aransas Pass	Ω	12.2	51	14.6	15	84	160	
	Aransas Pass	Q	12.2	51	16.4	15	4	167	
	S. of Corpus Christi		12.2	51	18.3	15	07	164	
	Pass								
25	Cedar Bayou	Ω	12.2	51	7.3	15	0		
	San Jose Island	Ω	12.2	51	14.6	15	7	161	ر ـ
	San Jose Island	Q	12.2	51	12.8	15	0		
	San Jose Island	Q	12.2	51	14.6	15	4	176	
25 Jun	San Jose Island	Q	12.2	51	7.3	15	0		
	Aransas Pass	Ω	12.2	51	16.4	15	0		
30	S. of Aransas Pass		12.2	51	9.1	15	184	169	
	S. of Corpus Christi		12.2	51	14.6	15	12	186	
	Pass								
Jul	ND								

PWD-RP-3400-340-1/91

Dispersal of this publication conforms with Texas State Documents Depository Law, and it is available at Texas State Publications Clearinghouse and Texas Depository Libraries.