BIRDS

— OF —

GALVESTON ISLAND S T A T E P A R K



A FIELD CHECKLIST

INTRODUCTION

nce little more than a wavering sliver of grass-carpeted sand, Galveston Island now favors the inland cities massed along the shores of the bay that carries its name. The inhospitable nature of a windswept barrier island has been supplanted by more "accommodating" (to humans, at least) beach homes, condominiums and curio shops. Only at Galveston Island State Park has a vestige of the original island coastal prairie been protected. This 2,000-acre park, situated six miles southwest of the western tip of Galveston's seawall, offers visitors a rare, intimate view of a barrier island ecosystem. Biological communities that once so typified this northernmost barrier island on the Texas coast still flourish at this vestigial spot as though the footsteps of Cabeza de Vaca and Jean Lafitte were only imagined imprints in the beach sand.

Spanning the width of the island from the Gulf of Mexico across to West Galveston Bay, the park is comprised of a mosaic of coastal habitats which hosts a surprising variety of birds. The secret to finding any specific bird is to search the right habitat at the right time of year. This checklist has been compiled in hopes that visitors to the park will have adequate information to locate the birds that are the object of their quest. To that end, pay specific attention to the habitat codes that accompany each species on the checklist. These classifications are based upon the general habitat types utilized by the birds in the parks and are not necessarily reflective of vegetational communities. The habitats that are of interest to birders are as follows:

Open Waters of the Gulf of Mexico

Extending from the beach, gulfward to the horizon, the stretch of the Gulf of Mexico that is visible from the park is noteworthy for the seabirds that periodically pass over and through its waters. In addition to the ubiquitous pelicans, gulls, and terns that congregate over the gulf, birders may also see Northern Gannets, Pomarine and Parasitic jaegers in the winter and during migrations and Magnificent Frigatebirds in summer and fall. There are winters when sea ducks gather offshore, presenting the rare opportunity to see Long-tailed Duck, and Black, Surf and White-winged scoters in Texas. In early spring, migrating Tricolored Herons, White Ibis, Fulvous Whistling-Ducks and Blue-winged Teal can be seen returning from their southerly wintering grounds in an endless stream moving north along the coast.

Sandy Beach/Dune

This littoral habitat ranges from the water line landward through the fore-island dunes. The vegetation on the gulfward dune faces is dominated by succulents and salt-tolerant species of plants such as gulf croton (Croton punctatus), sea-purslane (Sesuvium portulacastrum), fiddleleaf morningglory (Ipomoea stolonifera), bitter or seaside panicum (Panicum amarum) and camphor daisy (Heterotheca subaxillaris). This community grades into a seacoast bluestem (Schizachyrium scoparium) – gulfdune paspalum (Paspalum monostachyum) community on the landward dune slopes. Gulls and terns roost in mixed flocks along the open beaches, and unusual species such as Franklin's and Glaucous gulls and Common Tern are usually associated only with these groups. Several species of shore-birds, including the endangered Piping Plover, can be seen feeding in the wet sand along the water line. Horned Larks and several species of sparrows (winter) can be found foraging for seeds in the heavy grasses immediately behind the dunes. In the winter and during migrations, both Merlins and Peregrine Falcons perch on and along the beaches, waiting for an unsuspecting shorebird or migrant passerine to wander their direction.

Coastal Grassland

Approximately 600 acres of park property were originally midgrass grassland with seacoast bluestem and gulfdune paspalum as dominants. Until purchased by TPWD for the creation of a state park, this grassland was heavily grazed by cattle. After acquisition, this disturbed prairie left ungrazed and unburned, quickly began a succession to woody shrub. Baccharis (*Baccharis halimifolia*) and the exotic Chinese tallow (*Sapium sebiferum*) came to dominate these upland areas, and many of the grassland species of birds were replaced by those more typical of eastern woodlands. TPWD recently began a program to restore these critical coastal grasslands through prescribed fire and cyclical mowing. As these grasslands recover, many prairie-inhabiting birds will return to occupy this unique coastal habitat. For example, we should see birds such as Short-eared Owl and Le Conte's Sparrow become increasing common in the winter, with species such as the Bobolink appearing in the migration.

Coastal Scrub/Woodland

Before the arrival of Europeans, Galveston Island lacked significant tracts of native woodlands. Historical records indicate that woodland vegetation was limited to a few live oaks (*Quercus virginiana*) near the western tip of the island.

The trees and shrubs in the park, therefore, have become established since and as a direct result of the arrival of European man. Species such as eastern live oak, Hercules club (Zanthozylum clava-herculis), red mulberry (Morus rubra), wollybucket (Bumelia lanuginose), sugar hackberry (Celtis laevigata) and gulf black willow (Salix nigra) are native to the Texas coast and their presence in the park, although a result of man's activities, should be considered natural. Exotic species, such as Chinese tallow, slash pine (Pinus elliottii), salt cedar (Tamarix gallica), Japanese honevsuckle (Lonicera japonica) and chinaberry (Melia azedarach) originated either as ornamentals cultivated around the homes of early homesteaders or as windbreaks planted for cattle. These exotics are decidedly anomalous and are being eradicated from the park as a part of the program to restore the native ecosystem. Baccharis, although a native woody shrub, is being reduced from its position of dominance in these disturbed grasslands to a relative abundance more in line with its natural occurrence. Several live oak mottes, or tree islands, will be maintained within the park to enhance the native woodland habitat available for Neotropical migrants such as flycatchers, vireos, and warblers. The woodlands in concert with the restored coastal grasslands and associated wetlands will assure a diversity of habitats and bird species at Galveston Island State Park.

Freshwater Pond/Swale Assemblage

In general, fresh water is restricted and its sources are ephemeral on a barrier island—Galveston Island State Park is no exception. An impressive array of wetland birds, however, can be found in and around the few freshwater swales and ponds in the park. Many of these ponds and swales, particularly those immediately landward of the fore-island dunes, are lined with dense stands of common cattail (Typha domingensis) and common reed (Phragmites australis). These lagoons are a natural geological feature of the dune-upland interface along Texas' barrier islands, but have been largely destroyed on Galveston Island by draining and filling for development. This assemblage includes both the open waters of these ponds themselves and the freshwater marshes which border them. Several species of dabbling ducks winter in these shallow waters, and herons and egrets are always present feeding on the shores. Many migratory species of shorebirds only appear in the park to feed on the exposed mud around these ponds, and these birds are therefore more common in the park in years when a lack of rain has lowered freshwater levels. Also check the salt cedars around these ponds during spring and fall migration. Vireos, warblers, buntings, and orioles are attracted by the food and protection offered by these pseudo-riparian woodlands. During migrant groundings induced by inclement weather (particularly in the spring), these trees may appear to be festooned with the ornaments of a shattered avian rainbow.

Tidal Slough/Saltmarsh Assemblage

Unlike the narrow stretch of sandy beach and low dune fields on the seaward side of the island, the bay side is characterized by broad expanses of smooth cordgrass (Spartina alterniflora) marsh. Several sloughs and tidal bayous indent the bay margin, and these shallow tidal waters teem with a profusion of fishes and invertebrates. These wetlands are critical for the health of Galveston Bay. Juvenile fish and invertebrates are sheltered by the roots and stems of the grasses, and the marshes are a major source of detrital export for the bay system as a whole. The abundance of prey may attract hundreds of herons and egrets on a given day. Other long-legged waders such as Roseate Spoonbill and White Ibis can best be seen in this habitat. The distribution of several salt-marsh specialties, such as Clapper Rail and Seaside Sparrow, is restricted to these smooth cordgrass marshes. The muddy substrate exposed at low tide offers a rich feeding ground for many shorebirds such as Black-bellied Plover, Willet, Dunlin and Short-billed Dowitcher. Grebes, cormorants, and loons are attracted by the fish that crowd the sloughs in winter. Even diving ducks such as Bufflehead may at times venture into the deepest waters of Oak and Carancahua bayous during that season.

Emergent tidal marshes and transitional habitats are extremely elevation-sensitive. subsidence of the island over the past half-century has resulted in losses of up to 90% of these critical wetlands in West Galveston Bay marshes, including those at the State park. In the late 1990's a project was undertaken to protect and restore remaining tidal marshes at the State Park and to protect shallow bay bottom wetlands from further erosion and turbidity. The large fabric-covered "geo-tubes" that can be seen protecting the park's coves, and the rows of levees in shallow water of Carancahua Cove, now covered with smooth cordgrass, are the products of the project. In addition to increasing the production of small fishes, crustaceans, and other invertebrates, which serve as food for shore and wading birds, these structures act as islands for resting and nesting. In winter they attract species such as the Least Sandpiper, American Oystercatcher, and Marbled Godwit. In spring and summer, the levees attract

hundreds of Black Skimmers and other colonial nesting water birds.

The protected coves have also become home to expansive seagrass beds, which were extirpated from West Galveston Bay by turbidity and changing water quality in the 1960's. These beds attract a diversity of fishes and other wildlife, which in turn attract feeding birds. The grasses themselves, especially shoalgrass (*Halodule wrightii*), also provide food for wintering waterfowl such as Canvasbacks and Redheads.

Bay Margin/Tidal Mud Flat

At ebb tide, particularly during winter "northerns" when the bay waters are at their lowest, vast areas of normally inundated mudflats and oyster reefs are exposed to feeding shorebirds, herons, and egrets. The reefs may attract one or two American Oystercatchers and an extremely low tide is ordinarily the only opportunity to see this rarity in or near the park. Reddish Egrets will often prey upon the fish left stranded by the receding tides, and Marbled Godwits, normally absent in the park, will wade through the shallow waters looking for bristle worms. A distinctive tidal flat vegetation has evolved upon the saline soils found immediately inland of the bay margins. Plants such as glasswort (Salicornia europaea), saltwort (Batis maritime), saltgrass (Distichlis spicata) and sea ox-eye daisy (Borrichia frutescens) predominate. Drainage culverts and roofs provide nest sites for Barn Swallows. Purple Martins nest in houses erected specifically for that purpose. Killdeer and Common Nighthawks nest in small depressions (scrapes) in oyster shell and gravel roads in the park.

Aerial Feeder/Transient

Although rarely landing, there are several species of birds that are typically seen only when they fly over or through the park. Except for the Barn Swallows breeding within the park in spring, migrating swallows feed on the wing and therefore rarely stop within the park's boundaries. Occasionally, hawks are viewed riding thermals (kettling) high overhead, and flocks of shorebirds can be seen in spring flying in from the Gulf of Mexico after non-stop flights from Central and South America. Gulls and terns are forever crossing the park as they move between the gulf and bay, especially in the breeding season. Barn Owls, the only resident owl within the park, are most often seen searching for prey as they hover over the open coastal grasslands at twilight.

This bird list was prepared by Ted Eubanks, Jr. of Houston, edited and revised by Jim Stevenson of Galveston in 2001 and updated in 2018, 2020 and 2024.

Please help protect the natural avian communities in our parks by refraining from using playback tapes of bird songs. Frequent use of these tapes disrupts normal avian activity patterns, disrupts essential territorial behavior and may lead to nest failure. Thank you for your cooperation.

LEGEND

Habitat Codes

- 1 Open waters of the Gulf of Mexico
- 2 Sandy beach/dune
- 3 Coastal grasslands
- 4 Coastal scrub/woodland
- 5 Freshwater pond/swale assemblage
- 6 Tidal slough salt marsh assemblage
- 7 Bay margin/tidal mud flat
- 8 Open waters of West Galveston Bay
- 9 Rookeries/breeding colonies
- 10 Manmade structures
- 11 Aerial feeders/transient

Seasons

SP - Spring: March-May

SU - Summer: June-July

FA - Fall: August-November

WI - Winter: December-February

Status Codes (depending on season)

C - Common

U - Uncommon

0 – Occasional

R - Rare

X - Accidental

Status/Habitat Modifiers


- * Species documented as breeding in park
- ? Designated status and/or habitat uncertain
- M Habitat utilized primarily during migration
- [I] Irruptive species, with numbers fluctuating widely each year


CHECKLIST

	SP	SU	FA	WI	Habitat
Black-bellied Whistling-Duck	. С	С	0	0	5, 6
Fulvous Whistling-Duck	. 0	R			1M, 5
Greater White-fronted Goose			0	U	1M, 8, 11
Snow Goose			0	U	1M, 8, 11
Ross's Goose			R	R	1M, 8, 11
Gadwall	. С		C	C	5, 6, 8
American Wigeon	. С		C	C	5, 6, 8
Mallard	. R		R	0	5, 6, 8
Mottled Duck	. С	C*	C	C	5*, 6*
Wood Duck			R	0	5, 6, 8
Blue-winged Teal	. С	R*	C	0	5, 6, 8
Cinnamon Teal	. R		R	R	5
Northern Shoveler	. С		C	C	5, 6, 8
Northern Pintail	. U		U	U	1M, 5, 6, 8
Green-winged Teal	. U		U	U	1M, 5, 6, 8
Canvasback	. U		U	U	5, 6, 8
Redhead	. 0		0	0	5, 6, 8
Ring-necked Duck	. U		U	U	5, 6, 8
Greater Scaup	. R		R	U	1, 5, 6, 8
Lesser Scaup	. С	R	C	C	1, 5, 6, 8
Surf Scoter				R	1

	SP	SU	FA	WI	Habitat
White-winged Scoter				R	1
Black Scoter				R	1
Long-tailed Duck				R	1
Bufflehead	U			U	5, 6, 8
Common Goldeneye				0	1,8
Hooded Merganser	R			R	5, 6
Red-breasted Merganser	C		C	C	1, 6, 8
Ruddy Duck	U		U	U	5, 6
Pied-billed Grebe	C	0*	C	C	5*,6
Horned Grebe	R		R	R	1, 6, 8
Eared Grebe	U		U	U	1, 6, 8
Rock Pigeon	0	0*	0	0	10*, 11
Eurasian Collared-Dove	U	U*	U	U	10*, 11
Inca Dove	U	U*	U	U	4*
Common Ground-Dove	R		R	R	4
White-winged Dove	0	0*	C	0	4*
Mourning Dove	C	C*	C	C	4*
Black-billed Cuckoo	U		R		4
Yellow-billed Cuckoo	C	R	C		4
Groove-billed Ani	0	R	U	0	4
Lesser Nighthawk	0		0		4, 10, 11
Common Nighthawk	C	C*	C		4, 10*, 11
Chuck-will's-widow	C		C		4, 11
Eastern Whip-poor-will	0		0		4, 11
Chimney Swift	C		C		11
Buff-bellied Hummingbird	R			R	4
Ruby-throated Hummingbird	C	R	C		4
Black-chinned Hummingbird			R	R	4
Rufous Hummingbird			R	R	4

	SP	SU	FA	WI	Habitat
Broad-tailed hummingbird				R	4
King Rail	U	R	0	U	5, 6, 7
Yellow Rail	R		R	R?	5, 6
Black Rail	R	R*?			5, 6
Clapper Rail	C	C*	C	C	6*, 7
Virginia Rail	0		0	0	5, 6
Sora	U		U	0	5, 6
Purple Gallinule	0	R*	0		5*
Common Gallinule	U	0*	U	U	5*
American Coot	C	R	C	C	5
Sandhill Crane	0		0	0	3, 11
Black-necked Stilt	C	C*	C	R	5*, 6*
American Avocet	0		0	0	5, 6, 7
American Oystercatcher	R	R*	R	R	7, 9*
Black-bellied Plover	C	0	C	C	2, 6, 7
American Golden-Plover	C				3, 11
Snowy Plover	0		0	0	2,7
Wilson's Plover	C	C*	C	R	2, 7*
Semipalmated Plover	U	R	U	U	2, 5, 6, 7
Piping Plover	U		U	U	2, 7
Killdeer	C	C*	C	C	3, 5, 6, 10*
Upland Sandpiper	U		0		3, 11M
Whimbrel	U		U	R	3, 5, 6, 7
Long-billed Curlew	C		C	C	3, 5, 6, 7
Hudsonian Godwit	R				5, 11M
Marbled Godwit	U		U	U	6, 7
Ruddy Turnstone	C	0	C	C	2, 6, 7
Red Knot	U		U	0	2
Stilt Sandpiper	U		U	R	5, 6, 11M





		SP	SU	FA	WI	Habitat
	Sanderling	С	0	C	C	2, 6, 7
	Dunlin	C		C	C	5, 6, 7
	Baird's Sandpiper	0		0		2, 3, 5, 6, 7
	Least Sandpiper	C		C	C	5, 6, 7
	White-rumped Sandpiper	0				5, 6, 7, 11M
	Buff-breasted Sandpiper	R		R		3
	Pectoral Sandpiper	U		U		3, 5, 11M
	Semipalmated Sandpiper	U		0		5
	Western Sandpiper	C	R	C	C	2, 5, 6, 7
	Short-billed Dowitcher	C	0	C	C	6, 7
	Long-billed Dowitcher	U		U	U	5
	Wilson's Snipe	C		C	C	5
	Spotted Sandpiper	U		U	0	5, 6, 7
	Solitary Sandpiper	U		U		5
	Greater Yellowlegs	C		C	C	5, 6, 7, 11M
	Willet	C	C*	C	C	2, 5, 6*, 7*
	Lesser Yellowlegs	C		C	U	5, 6, 11M
	Wilson's Phalarope	0		0		5, 6, 7
	Pomarine Jaeger	R		R		1M, 8, 11
	Laughing Gull	C	C*	C	C	1, 2, 7, 8, 9*, 11
	Franklin's Gull	R		U		1, 2, 8, 11M
	Bonaparte's Gull	C		C	C	1, 2, 8
	Ring-billed Gull	C	0	C	C	1, 2, 8, 11
	Herring Gull	C	0	C	С	1, 2, 8, 11
	California Gull	0				1, 2, 8
	Gull-billed Tern	U	U*	U	0	5, 6, 9*, 11
	Caspian Tern	C	C*	С	C	1, 2, 5, 6, 8, 9*, 11
	Royal Tern	C	C*	С	C	1, 2, 5, 6, 8, 9*, 11
	Sandwich Tern	C	C*	С	R	1, 2, 5, 6, 8, 9*, 11
_						

	SP	SU	FA	WI	Habitat
 Common Tern	U	0	U	R	1, 2
 Forster's Tern	C	C*	C	C	1, 2, 5, 6, 8, 9*, 11
 Least Tern	C	C*	C		1, 2, 5, 6, 8, 9*, 11
 Black Tern	C	U	C		1, 2, 5, 6, 8, 11
 Black Skimmer	C	C*	C	C	1, 2, 5, 6, 8, 9*, 11
 Pacific Loon	X				
 Common Loon	U	R	0	U	1, 8
 Wood Stork		R	R		5, 6
 Magnificent Frigatebird	0	U	U		1, 8, 11
 Northern Gannet	U		0	U	1
 Neotropic Cormorant	C	C	C	0	6, 8
 Double-crested Cormorant	C		C	C	1, 6, 8
 Anhinga	R		R		5, 11M
 American White Pelican	C	R	C	C	1, 6, 8
 Brown Pelican	0	U	U	0	6, 8
 American Bittern	U		U	U	5, 6
 Least Bittern	U	U*	0		5*, 6*
 Great Blue Heron	C	C*	C	C	5, 6, 7, 9*
 Great Egret	C	C*	C	C	5, 6, 7, 9*
 Snowy Egret	C	C*	C	C	5, 6, 7, 9*
 Little Blue Heron	C	U	C	0	5, 6, 7
 Tricolored Heron	C	C*	C	U	5, 6, 7, 9*
 Reddish Egret	C	C*	C	0	6, 7, 9*
 Cattle Egret	C	C*	C	U	3, 5, 7, 9*
 Green Heron	C	C*	C	R	5*, 6
 Black-crowned Night-Heron	C	C*	C	C	5, 6, 7, 9*
 Yellow-crowned Night-Heron	U	0	U	R	5, 6, 7
 White Ibis	C	C	C	C	5, 6, 7
 Glossy Ibis	R				5

		SP	SU	FA	WI	Habitat
White-faced Ibi	is	U	U	U	U	5, 6
Roseate Spoonl	bill	C	C*	C	U	5, 6, 7, 9*
Black Vulture		U	U	U	U	10, 11
Turkey Vulture		U	U	U	U	10, 11
Osprey		U	R	U	0	1, 5, 6, 8, 11M
Swallow-tailed	Kite	R				11M
White-tailed Ki	te	U	0*	U	U	3, 4*
Mississippi Kite		0		0		4, 11M
Northern Harri	er	C	R*	C	C	3*
Sharp-shinned	Hawk	U		U	0	4, 11M
Cooper's Hawk		U		U	0	4, 11M
Red-shouldered	d Hawk	U		U	U	4, 10, 11
Broad-winged	Hawk	0		U		4, 11M
Red-tailed Haw	/k	U		U	U	3, 4, 11
Barn Owl		C	C*	C	C	3, 4*, 10*
Great Horned ()wl	U		U	U	3, 4
Short-eared Ov	vl	U		U	U	3, 4[I]
Belted Kingfish	er	C	R	C	C	5, 6
Red-bellied Wo	odpecker	U		U	U	4
Yellow-bellied S	Sapsucker	U		U	U	4
Downy Woodpe	ecker	U	U	U	U	4
Northern Flick	er	U		U	U	4
Crested Caraca	ra	U	0*	U	U	3, 4*
American Kestr	el	C		C	C	3, 4, 11M
Merlin		U		U	0	2, 4, 11M
Peregrine Falco	on	U		U	0	2, 11M
Olive-sided Flyo	catcher	0		U		4
Eastern Wood-	Pewee	C		C		4
Yellow-bellied I	Flycatcher	0		U		4

	SP	SU	FA	WI	Habitat
Acadian Flycatcher	U		U		4
Alder Flycatcher			U		4
Willow Flycatcher			0?		4
Least Flycatcher			C	R	4
Eastern Phoebe			C	C	4
Vermilion Flycatcher	R		R	R	4
Ash-throated Flycatcher	R		R		4
Great Crested Flycatcher	С	R	C		4
Tropical Kingbird			R		4
Western Kingbird	U		U		4
Eastern Kingbird	С	C*	C		4
Scissor-tailed Flycatcher	С	C*	C		4
Fork-tailed Flycatcher				0	4
Loggerhead Shrike	С	C*	C	C	4*
White-eyed Vireo	С		C	R	4
Blue-headed Vireo	U		U	U	4
Yellow-throated Vireo	U		U		4
Warbling Vireo	0		U		4
Philadelphia Vireo	U		0		4
Red-eyed Vireo	С		C		4
Blue Jay	0	0	0	0	4[I]
Carolina Chickadee	0	R	R	U	4
Horned Lark	U	U^*	U	U	2*, 7*
Purple Martin	С	C*	C		10*, 11
Tree Swallow	С		C	0	11
No. Rough-winged Swallow	С		C		11
Bank Swallow	С		C		11
Cliff Swallow	0		0		11
Cave Swallow	U	R	U	R	11

	SP	SU	FA	WI	Habitat
 Barn Swallow	С	C*	С		11
 Carolina Wren	R	R	R	R	4
 House Wren	C		C	C	4
 Winter Wren			R		4
 Sedge Wren	C		C	C	3
 Marsh Wren	U	U*	U	U	5*
 Blue-gray Gnatcatcher	C		C	U	4
 Golden-crowned Kinglet	U		U	U	4[I]
 Ruby-crowned Kinglet	C		C	C	4
 Veery	U		R		4
 Gray-cheeked Thrush	U		R		4
 Swainson's Thrush	C		C		4
 Hermit Thrush	U		U	U	4
 Wood Thrush	C		C		4
 American Robin	C		C	C	4
 Gray Catbird	C		C	0	4
 Brown Thrasher	U		U	U	4
 Northern Mockingbird	C	C*	C	C	4
 Eastern Bluebird	U			0	4
 European Starling	C	C*	C	C	4, 10*
 Cedar Waxwing	U		U	U	4[I]
 Scaly-breasted Munia			R		4
 House Sparrow	0	0*	0	0	10*
 American Pipit	U		U	U	3, 5
 Sprague's Pipit	R		R	R	3
 House Finch	0		0	0	4[I]
 American Goldfinch	U		U	U	4
 Eastern Towhee			U	U	4[I]
 Chipping Sparrow	0		0	0	4

	SP	SU	FA	WI	Habitat
Clay-colored Sparrow			R		3
Field Sparrow	. 0		0	0	3, 4
Vesper Sparrow	. 0		0	0	3, 4
Lark Sparrow	. 0		0		4
Savannah Sparrow	. С		C	C	3
Grasshopper Sparrow	. 0		0	0	3
LeConte's Sparrow	. 0		0	0	3
Nelson's Sparrow	. U		U	U	6
Seaside Sparrow	. С	C*	C	C	6*
Song Sparrow	. U		U	U	3, 4
Lincoln's Sparrow	. С		C	C	3, 4
Swamp Sparrow	. С		C	C	5, 6
White-throated Sparrow	. U		U	U	4
White-crowned Sparrow	. 0		0	0	4[I]
Harris's Sparrow			0	0	4[I]
Dark-eyed Junco	. R		R	R	4[I]
Yellow-breasted Chat	. U		U		4
Bobolink	. R				3[I]
Red-winged Blackbird	. С	C*	C	C	3*, 4*, 5, 6
Eastern Meadowlark	. С	C*	C	C	3*
Western Meadowlark	. R?		R?	R?	3
Yellow-headed Blackbird	. R		R		3, 5
Common Grackle	. U	U	U	U	3, 4, 10
Boat-tailed Grackle	. U	U	U	U	5, 6, 10
Great-tailed Grackle	. С	C*	C	C	4*, 10*
Bronzed Cowbird	. R	R*?	0		3, 4
Brown-headed Cowbird	. С	C*	C	C	3, 4*
Brewer's Blackbird	. R			R	2, 3
Orchard Oriole	. С	C*	C		4*

		SP	SU	FA	WI	Habitat
Ba	altimore Oriole	С		С		4
Bl	lue-winged Warbler	C		C		4
Go	olden-winged Warbler	U		0		4
Te	ennessee Warbler	C		C		4
0:	range-crowned Warbler	C		C	C	4
Na	ashville Warbler	0		U		4
No	orthern Parula	C		C		4
Ye	ellow Warbler	C		C		4
Cl	hestnut-sided Warbler	C		C		4
M	agnolia Warbler	C		C		4
Ca	ape May Warbler	R				4
Bl	lack-throated Blue Warbler	R		R		4
Ye	ellow-rumped Warbler	C		C	C	4
Bl	lack-throated Green Warbler	C		C		4
Bl	lackburnian Warbler	C		C		4
Ye	ellow-throated Warbler	U		U		4
Pi	ine Warbler	R		R	R	4[I]
Pı	rairie Warbler	R		U		4
Bl	lack-throated Gray Warbler			0		4
Pa	alm Warbler	U		U	U	4
Ba	ay-breasted Warbler	C		C		4
Bl	lackpoll Warbler	U		R		4
Ce	erulean Warbler	U		R		4
Bl	lack-and-white Warbler	C		C		4
Ar	merican Redstart	C		C		4
Pı	rothonotary Warbler	C		C		4
W	orm-eating Warbler	C		C		4
Sv	wainson's Warbler	0		R		4
0	venbird	C		C		4

	SP	SU	FA	WI	Habitat
Northern Waterthrush	С		С		4, 5
 Louisiana Waterthrush	U		U		4, 5
 Kentucky Warbler	C		C		4
 MacGillvray's Warbler	R		R		4
 Mourning Warbler	R		U		4
 Common Yellowthroat	C	C*	C	C	3*, 4*, 5*
 Hooded Warbler	C		C		4
 Wilson's Warbler	0		U	R	4
 Canada Warbler	U		C		4
 Summer Tanager	C		C	R	4
 Scarlet Tanager	C		R		4
 Western Tanager	R		R	R	4[I]
 Northern Cardinal	U	U^*	U	U	4*
 Rose-breasted Grosbeak	C		C		4
 Blue Grosbeak	C		C		4
 Indigo Bunting	C		C	R	4
 Painted Bunting	C	0*	C		4*
 Dickcissel	C	0*	C		3*, 11M

The following species have been recorded in Galveston Island State Park, but should not be expected every year: Horned, Least and Western grebes, Masked Booby, Glossy Ibis, Wood Stork, Black Vulture, Black, Surf and White-winged scoters, Swallow-tailed Kite, Black Rail (possibly breeds), Eskimo Curlew (extinct?), Hudsonian Godwit, American Woodcock, Parasitic Jaeger, Lesser Black-backed Gull, Sooty Tern, Great Horned Owl, Buff-bellied and Rufous hummingbirds, Downy and Red-headed woodpeckers, Vermillion, Ash-throated and Brown-crested flycatchers, Say's Phoebe, Couch's Kingbird, Red-breasted Nuthatch, Eastern, Spotted and Green-tailed towhee, Dark-eyed Junco, Pyrrhuloxia, Western Meadowlark, Yellow-headed Blackbird and Purple Finch.

CITATION

Stevenson, J.M. March 2024. Birds of Galveston Island State Park: a field checklist. Natural Resource Program, Texas Parks and Wildlife, Austin, Texas.

In 1983, the Texas Legislature created the Special Nongame and Endangered Species Conservation Fund. This fund may be used for nongame wildlife and endangered species research and conservation, habitat acquisition and development and dissemination of information pertaining to nongame management. Money for this fund is obtained through private donations and sale of nongame wildlife art prints and stamps. This fund now gives Texans a unique opportunity to help support this state's valuable and sensitive nongame resources. Your individual contributions and purchases of nongame art prints and stamps will help determine the level of nongame conservation activities in Texas. For more information call toll-free (1-800-792-1112) or contact: Nongame and Endangered Species Program, Texas Parks and Wildlife, 4200 Smith School Road, Austin, Texas 78744.

Cover: Illustration of Clapper Rail by Clemente Guzman.

TPWD receives funds from DHS and USFWS. TPWD prohibits discrimination based on race, color, religion, national origin (including limited English proficiency), disability, age, and gender, pursuant to state and federal law. If you believe you have been discriminated against by TPWD, visit tpwd.texas. gov/nondiscrimination or call (512) 389-4800 for information on filing a complaint. To obtain information an alternative format, contact TPWD through Relay Texas at 7-1-1 or (800) 735-2980, or by email at accessibility@tpwd.texas.gov. (If you speak a language other than English and need assistance, email lep@tpwd.texas.gov. You can also contact Department of the Interior Office of Diversity, Inclusion, and Civil Rights, 1849 C Street, N.W., Washington, D.C. 20240, and/or U.S. Department of Homeland Security Office for Civil Rights and Civil Liberties (CRCL), Mail Stop #0190 2707, Martin Luther King, Jr. Ave., S.E. Washington, D.C. 20528.



Natural Resources Program Texas Parks and Wildlife 4200 Smith School Road Austin, Texas 78744